

GOSSIP COVERAGE CONTROL FOR ROBOTIC NETWORKS: DYNAMICAL SYSTEMS ON THE SPACE OF PARTITIONS*

FRANCESCO BULLO[†], RUGGERO CARLI[†], AND PAOLO FRASCA[‡]

Abstract. Future applications in environmental monitoring, delivery of services and transportation of goods motivate the study of deployment and partitioning tasks for groups of autonomous mobile agents. These tasks may be achieved by recent coverage algorithms, based upon the classic methods by Lloyd. These algorithms however rely upon critical requirements on the communication network: information is exchanged synchronously among all agents and long-range communication is sometimes required. This work proposes novel coverage algorithms that require only *gossip communication*, i.e., asynchronous, pairwise, and possibly unreliable communication. Which robot pair communicates at any given time may be selected deterministically or randomly. A key innovative idea is describing coverage algorithms for robot deployment and environment partitioning as dynamical systems on a space of partitions. In other words, we study the evolution of the regions assigned to each agent rather than the evolution of the agents' positions. The proposed gossip algorithms are shown to converge to centroidal Voronoi partitions under mild technical conditions.

Our treatment features a broad variety of results in topology, analysis and geometry. First, we establish the compactness of a suitable space of partitions with respect to the symmetric difference metric. Second, with respect to this metric, we establish the continuity of various geometric maps, including the Voronoi diagram as a function of its generators, the location of a centroid as a function of a set, and the widely-known multicenter function studied in facility location problems. Third, we prove two convergence theorems for dynamical systems on metric spaces described by deterministic and stochastic switches.

Key words. Cooperative control, multiagent systems, gossip communication, geometric optimization, centroidal Voronoi tessellations, Lloyd algorithm

AMS subject classifications. 37N35, 68T40, 68W15, 93D20, 03H05

1. Introduction. In the not too distant future, networks of coordinated autonomous robots will perform a broad range of environmental monitoring and logistic tasks. Robotic camera networks will monitor airports and other public infrastructures. Teams of vehicles will perform surveillance, exploration and search and rescue operations. Groups of robots will enable novel logistic capacities in the transportation of goods and the delivery of services and resources to customers. New applications will be enabled by the ongoing decreases in size and cost and the increases in performance of sensors, actuators, communication devices and computing elements.

In these future applications, load balancing algorithms will dictate how the workload is shared and assigned to the individual robots. In other words, robotic resources will be assigned and deployed to competing requests in such a way as to optimize some performance metric. Remarkably, load balancing problems in robotic networks are often equivalent to robotic deployment and environment partitioning problems. For example, in surveillance applications, optimal sensor coverage is often achieved by

*To appear in the SIAM Journal on Control and Optimization. Submitted on August 23, 2010. This work was supported in part NSF grant IIS-0904501 and by ARO MURI grant W911NF-05-1-0219 and ONR grant N00014-07-1-0721. A preliminary and incomplete version of this work appeared in the Proceedings of the 2009 American Control Conference, pages 2228-2235, St. Louis, Missouri, USA. The authors gratefully acknowledge Prof. Andrew R. Teel for Remark 4.4 and Prof. Ramon van Handel for his remarks on the set of closed subsets and the symmetric distance metric.

[†]Francesco Bullo and Ruggero Carli are with the Center for Control, Dynamical Systems and Computation, University of California at Santa Barbara, Santa Barbara, CA 93106, USA, {bullo, carlirug}@engineering.ucsb.edu.

[‡]Paolo Frasca is with the Dipartimento di Matematica, Politecnico di Torino, paolo.frasca@polito.it.

partitioning the environment and assigning individual robotic sensors to individual regions of responsibility. Similarly, in the transportation of goods or delivery of services, minimizing the customer wait-time is equivalent to a multi-vehicle routing problem and, in turn, to computing optimal depot positions and regions of responsibility.

Motivated by these scenarios, this paper considers the two following interrelated problems. The *deployment problem* for a robotic network amounts to the design of coordination algorithms that lead the robots to be optimally placed in an environment of interest. Deployment performance is characterized by an appropriate network utility function that measures the deployment quality of a given configuration. The *partitioning problem* is the design of coordination algorithms that lead the robots to optimally partition the environment into subregions of interest; even here the objective is usually achieved through the design of appropriate utility functions.

Literature review. The “centering and partitioning” algorithm originally proposed by Lloyd [26] and elegantly reviewed in the survey [15] is a classic approach to facility location and environment partitioning problems. The Lloyd algorithm computes centroidal Voronoi partitions as optimal configurations of an important class of objective functions, called multicenter functions. Besides their intended application to quantization theory [19], centroidal Voronoi partitions have widespread applications in numerous disciplines, including statistical pattern recognition [23], geometric optimization [3] and spatial resource allocation [13]. Recent mathematical interest has focused on convergence analysis [14], bifurcation analysis of low dimensional problems [41], and anisotropic partitions [16], among other topics.

Distributed and robotic versions of the Lloyd algorithm have been recently developed in the multiagent literature; see the text [10, Chapter 5 and literature notes in Section 5.4]. We briefly review this growing literature in what follows. Generalized centroidal Voronoi partitions are shown in [18] to be asymptotically optimal for estimation of stochastic spatial fields by sensor networks. Boundary coverage problems and convergence rates are studied in [38] and [29], respectively. Convergence to centroidal Voronoi partitions is established in [4] for a class of communication-less sensor-based algorithms (related to the classic clustering work by MacQueen [28]). In [37] adaptive coverage controls are proposed for environments described by unknown density functions. In [32] partitioning policies are shown to achieve optimal load balancing in vehicle routing problems, i.e., problems in which a robotic network provides service to customers that arrive in real time in the environment. Finally, some anisotropic coverage problems and experimental implementation aspects are treated in [21].

Territory partitioning via competitive behaviors is a classic topic in behavioral ecology; see the comprehensive survey [1]. For example, it is known [2] that the foraging behavior of conflicting colonies of red harvester ants (*Pogonomyrmex barbatus*) results in non-overlapping dominance regions that resemble Voronoi partitions. Non-overlapping dominance regions akin to centroidal Voronoi partitions are documented in [6, 39, 15] for the mouthbreeder fish (*Tilapia mossambica*). Territoriality behavior and competition among prides of African lions (*Panthera leo*) are discussed in [30]. Overall, numerous animal species achieve territory partitioning without a central coordinating entity and without synchronized communication, but rather relying upon asynchronous accidental interactions and stigmergy. To the best of our knowledge, asynchronous territory partitioning has been barely studied, see [39] for introductory ideas about animal behavior, and mathematical models and analysis are lacking.

To finalize the literature review, here is a synopsis of other mathematical ideas that we bring to bear on deployment and partitioning problems. First, we adopt

the so-called *gossip communication model*, in which only peer-to-peer asynchronous communication links are required. This communication model is widely studied in the wireless communication literature; example references include [24, 7]. Moreover, we consider control systems on a non-Euclidean state space; the interest for non-Euclidean spaces has a rich history in nonlinear control theory, dynamical systems and robotics, including the early work [8] and a recent application to multiagent systems [36]. Finally, we adopt various tools from topology and from the study of hyperspaces of sets [31].

Statement of contributions. This paper uncovers novel mathematical principles and tools of relevance in coordination problems and multiagent systems. We tackle partitioning and coverage control algorithms in innovative ways. First, we design algorithms that require only gossip communication, i.e., asynchronous, pairwise, and possibly unreliable communication. Gossip communication is a simple, robust and effective protocol for noisy and uncertain wireless environments. Gossip communication may be implemented in wandering robots with short-range unreliable communication (an illustrative motion coordination strategy is reported in the report [9]). Second, we propose a change of perspective in coverage control and multicenter optimization. Classically [10, 18, 11, 32, 37, 41], the state space for the coverage algorithms are the agents' positions, i.e., as a function of the agents' positions the environment is divided into regions and regions are assigned to each agent. Note that in this classical approach, every movement of an agent is reflected in a change of both its own assigned region and its neighboring regions. Clearly, this rigidity conflicts with allowing unreliable and asynchronous communication. Instead, in our approach, the agents' positions are no longer a concern: the state space is the space of partitions of the environment and the algorithm dictates how to update the regions. As the space of partitions is much richer than the space of the agents' positions, we gain more freedom in the design of partition optimization algorithms, and in particular the possibility to use gossip communication.

Within the innovative context of gossip communication and partition-based mechanisms, we devise a novel algorithm for multicenter and coverage optimization. Our gossip coverage algorithm is a peer-to-peer version of Lloyd algorithm and aims to compute centroidal Voronoi partitions. Which robot pair communicates at any given time is the outcome of either a deterministic or a stochastic process. We also propose a modified version that restricts communication exchanges to adjacent regions and that has suitable continuity properties. Simulations illustrate that our algorithms successfully compute centroidal Voronoi partitions.

To formally establish the convergence properties of our proposed gossip coverage algorithms, we perform a detailed mathematical analysis composed of three steps. First, we develop suitable versions of the Krasovskii-LaSalle invariance principle for dynamical systems on metric spaces described by deterministic and stochastic switches. Convergence to a set of fixed points is achieved under a uniform deterministic or stochastic persistency condition. Second, we establish the continuity of various geometric maps, including (1) the Voronoi diagram as a function of its generators, (2) the location of the generalized centroid as a function of a set, (3) the widely-known multicenter function studied in facility location, and (4) the gossip coverage algorithms. These continuity properties are established with respect to the symmetric distance metric in the space of partitions. Third, we study the topology of the space of partitions. With respect to the symmetric difference metric we prove the compactness of a relevant subset of partitions. Specifically, we focus on partitions

whose component regions are the union of a bounded number of convex sets.

In summary, relying upon our extensions of the invariance principle, the compactness of a subset of the set of partitions, and the continuity of the various relevant maps, we establish the convergence properties of the proposed gossip algorithms. In short, the algorithms converge to the set of centroidal Voronoi partitions under mild technical assumptions and under the assumption that the gossip communication exchanges satisfy either a deterministic or a stochastic persistency condition.

The interest for asynchronous state updates is not new in the coverage control literature. The asynchronous algorithms in [11] rely upon the theory of deterministic and stochastic gradient algorithms presented in [40]. However, such stochastic gradient algorithms entail a significant limitation: convergence can be shown only under the assumption that the update step be small enough and no estimate of its size is available. On the contrary, the partitions-based framework presented here has the following advantages: there are no limitations on the step sizes, an explicit Lyapunov function is known, and the approach can be easily extended to discrete and more general setups; see [17] for some preliminary work in this direction.

Organization and notations. The paper is structured as follows. In Section 2 we review multicenter optimization and coverage control ideas. In Section 3 we state our asynchronous territory partitioning problem, provide a solution via the gossip coverage algorithm, state the convergence properties of the algorithm and report some simulation results. The following Sections 4, 5 and 6 develop the mathematical machinery required to prove the convergence results. Section 4 contains the convergence theorems extending the Krasovskii-LaSalle invariance principle. Section 5 contains a discussion about the compactness properties of the space of partitions. Section 6 states the continuity properties of the relevant maps and functions and contains the proof of the main convergence results. Concluding remarks are given in Section 7.

We let $\mathbb{R}_{>0}$ and $\mathbb{R}_{\geq 0}$ denote the set of positive and non-negative real numbers, respectively, and $\mathbb{Z}_{\geq 0}$ denote the set of non-negative integer numbers. Given $A \subset \mathbb{R}^d$, we let $\text{int}(A)$, \bar{A} , ∂A and $\text{diam}(A)$ denote its interior, its closure, its boundary and its diameter, respectively. Given two sets X and Y , a *set-valued map* $T : X \rightrightarrows Y$ associates to an element of X a subset of Y .

2. A review of multicenter optimization and distributed coverage control. In this section we review a variety of known results in geometric optimization and in robotic coordination. In Subsection 2.1 we review the notion of environment partitions and we introduce the multicenter function as a way to define optimal environment partitions and optimal robot or sensor positions in the environment. In Subsection 2.2 we review some distributed control algorithms for agent motion coordination and environment partitioning based on the classic Lloyd algorithm.

2.1. Partitions, centroids and multicenter optimization. We let Q denote an environment of interest to be apportioned. We assume Q is a compact convex subset of \mathbb{R}^d with non-empty interior. Partitions of Q are defined as follows.

DEFINITION 2.1 (Partition). *An N -partition of Q , denoted by $v = (v_i)_{i=1}^N$, is an ordered collection of N subsets of Q with the following properties:*

- (i) $\cup_{i \in \{1, \dots, N\}} v_i = Q$;
- (ii) $\text{int}(v_i) \cap \text{int}(v_j)$ is empty for all $i, j \in \{1, \dots, N\}$ with $i \neq j$; and
- (iii) each set v_i , $i \in \{1, \dots, N\}$, is closed and has non-empty interior.

The set of N -partitions of Q is denoted by \mathcal{V}_N .

Let $p = (p_1, \dots, p_N) \in Q^N$ denote the position of N agents in the environment Q . Given a group of N agents and an N -partition, each agent is naturally in one-to-one correspondence with a component of the partition; specifically we refer to v_i as the *dominance region* of agent $i \in \{1, \dots, N\}$.

On Q , we define a *density function* to be a bounded measurable positive function $\phi : Q \rightarrow \mathbb{R}_{>0}$ and a *performance function* to be a locally Lipschitz, monotone increasing and convex function $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$. With these notions, we define the *multicenter function* $\mathcal{H}_{\text{multicenter}} : \mathcal{V}_N \times Q^N \rightarrow \mathbb{R}_{\geq 0}$ by

$$(2.1) \quad \mathcal{H}_{\text{multicenter}}(v, p) = \sum_{i=1}^N \int_{v_i} f(\|p_i - q\|) \phi(q) dq.$$

This function is well-defined because closed sets are measurable. We aim to minimize $\mathcal{H}_{\text{multicenter}}$ with respect to both the partition v and the locations p .

REMARKS 2.2 (Locational optimization). *As discussed in the introduction and in the survey [15], the multicenter function $\mathcal{H}_{\text{multicenter}}$ has numerous interpretations. Here we review two applications entailing robotic networks. First, in a vehicle routing and service delivery example [32], given vehicles at locations p_i , assume that $f(\|p_i - q\|)$ is the cost incurred by agent i to travel to service an event taking place at point q . Events take place inside Q with likelihood ϕ . Accordingly, $\mathcal{H}_{\text{multicenter}}$ quantifies the expected wait-time between event arrivals and agents servicing them.*

Second, in an environmental monitoring application [11], assume the robots aim to detect acoustic signals that originate and propagate isotropically in the environment. Because of noise and loss of resolution, the ability to detect a sound source originating at a point q from a sensor at position p_i is proportional to the signal-to-noise ratio (which degrades with $\|q - p_i\|$). If the performance function f equals minus the signal-to-noise ratio, then $\mathcal{H}_{\text{multicenter}}$ quantifies the expected signal-to-noise ratio and detection capacity for acoustic signals generated at random locations. \square

Among all possible ways of partitioning a subset of \mathbb{R}^d , one is worth of special attention. Define the *set of partly coincident locations* $S_N = \{p \in Q^N \mid p_i = p_j \text{ for some } i, j \in \{1, \dots, N\}, i \neq j\}$. Given $p \in Q^N \setminus S_N$, the *Voronoi partition of Q generated by p* , denoted by $V(p)$, is the ordered collection of the *Voronoi regions* $(V_i(p))_{i=1}^N$, defined by

$$(2.2) \quad V_i(p) = \{q \in Q \mid \|q - p_i\| \leq \|q - p_j\| \text{ for all } j \neq i\}.$$

In other words, the Voronoi partition is a map $V : (Q^N \setminus S_N) \rightarrow \mathcal{V}_N$. The regions $V_i(p)$, $i \in \{1, \dots, N\}$, are convex and, if Q is a polytope, they are polytopes. Now, given two distinct points q_1 and q_2 in \mathbb{R}^d , define the $(q_1; q_2)$ -*bisector half-space* by

$$(2.3) \quad H_{\text{bisector}}(q_1; q_2) = \{q \in \mathbb{R}^d \mid \|q - q_1\| \leq \|q - q_2\|\}.$$

In other words, the set $H_{\text{bisector}}(q_1; q_2)$ is the closed half-space containing q_1 whose boundary is the hyperplane bisecting the segment from q_1 to q_2 . Note that bisector subspaces satisfy $H_{\text{bisector}}(q_1; q_2) \neq H_{\text{bisector}}(q_2; q_1)$ and that Voronoi partition of Q satisfies $V_i(p_1, \dots, p_N) = Q \cap (\bigcap_{j \neq i} H_{\text{bisector}}(p_i; p_j))$.

Each region equipped with a density function possesses a point with a special relationship with the multicenter function. Define the scalar *1-center function* \mathcal{H}_1 by

$$(2.4) \quad \mathcal{H}_1(p; A) = \int_A f(\|p - q\|) \phi(q) dq,$$

where p is any point in Q and A is a compact subset of Q . Under the stated assumptions on the performance function f , the function $p \mapsto \mathcal{H}_1(p; A)$ is strictly convex in p , for any set A with positive measure (we postpone the proof to Lemma 6.1). Therefore, the function $p \mapsto \mathcal{H}_1(p; A)$ has a unique minimum in the compact and convex set Q . We define the *generalized centroid* of a compact set $A \subset Q$ with positive measure by

$$(2.5) \quad \text{Cd}(A) = \operatorname{argmin}\{\mathcal{H}_1(p; A) \mid p \in Q\}.$$

In what follows, it is convenient to drop the word ‘‘generalized,’’ and to denote by $\text{Cd}(v) = (\text{Cd}(v_1), \dots, \text{Cd}(v_N)) \in Q^N$ the vector of regions centroids corresponding to a partition $v \in \mathcal{V}_N$.

REMARK 2.3 (Quadratic and linear performance functions). *If the performance function is $f(x) = x^2$, then the global minimum of \mathcal{H}_1 is the centroid (also called the center of mass) of A , defined by*

$$\text{Cd}(A) = \left(\int_A \phi(q) dq \right)^{-1} \int_A q \phi(q) dq.$$

If the performance function is $f(x) = x$, then the global minimum of \mathcal{H}_1 is the median (also called the Fermat–Weber center) of A . See [10, Chapter 2] for more details. \square

Voronoi partitions and centroids have useful optimality properties stated in the following proposition and illustrated in Fig. 2.1.

PROPOSITION 2.4 (Properties of $\mathcal{H}_{\text{multicenter}}$). *For any partition $v \in \mathcal{V}_N$ and any point set $p \in Q^N \setminus S_N$,*

$$(2.6) \quad \mathcal{H}_{\text{multicenter}}(V(p), p) \leq \mathcal{H}_{\text{multicenter}}(v, p),$$

$$(2.7) \quad \mathcal{H}_{\text{multicenter}}(v, \text{Cd}(v)) \leq \mathcal{H}_{\text{multicenter}}(v, p).$$

Furthermore, inequality (2.6) is strict if any entry of $V(p)$ differs from the corresponding entry of v by a set with positive measure, and inequality (2.7) is strict if $\text{Cd}(v)$ differs from p .




FIG. 2.1. *Illustration of Proposition 2.4. The left figure shows a sample 2-partition v and point set p in a uniform square environment. The value of the cost function $\mathcal{H}_{\text{multicenter}}$ at (v, p) is diminished by either replacing v with the Voronoi partition generated by p (see central figure), or replacing p with the centroids of v (see right figure).*

The statements in Proposition 2.4 originate in the early work by S. P. Lloyd [26]; modern treatments are given in [15] and [10, Propositions 2.14 and 2.15]. Proposition 2.4 implies the following necessary condition: if a pair (v, p) with $p \notin S_N$ minimizes $\mathcal{H}_{\text{multicenter}}$, then $p = \text{Cd}(v)$ and $v = V(p)$ up to a set of measure zero. Accordingly, the partitions that minimize $\mathcal{H}_{\text{multicenter}}$ have the following property.

DEFINITION 2.5. *The partition $v \in \mathcal{V}_N$ is centroidal Voronoi if it has distinct centroids, that is, $\text{Cd}(v_i) \neq \text{Cd}(v_j)$ for all $j \neq i$, and $v = V(\text{Cd}(v))$.*

2.2. From Lloyd algorithm to distributed coverage control. Here we consider a group of robotic agents with motion, communication and computation capacities and we review a coverage control algorithm that determines the motion of each robot in a group and the associated partition in such a way as to minimize $\mathcal{H}_{\text{multicenter}}$. In what follows, we restrict our attention to $d = 2$, that is, we assume $Q \subset \mathbb{R}^2$.

To explain in what sense our algorithms are distributed, we introduce a useful graph. The *Delaunay graph* [12, 10] associated to the distinct positions $p \in Q^N \setminus S_N$ is the undirected graph with node set $\{p_i\}_{i=1}^N$ and with the following edges: (p_i, p_j) is an edge if and only if $V_i(p) \cap V_j(p)$ is non-empty.

The *coverage algorithm* we consider is a distributed version of the classic Lloyd algorithm [15] based on “centering and partitioning” for the computation of centroidal Voronoi partitions. The algorithm is distributed in the sense that each robot determines its region of responsibility and its motion plan based upon communication with only some neighbors. Specifically, communications among the robots takes place along the edges of the Delaunay graph. The distributed coverage algorithm is described as follows. At each discrete time instant $t \in \mathbb{Z}_{\geq 0}$, each agent i performs the following tasks: (i) it transmits its position and receives the positions of its neighbors in the Delaunay graph; (ii) it computes its Voronoi region with the information received; (iii) it moves to the centroid of its Voronoi region. In mathematical terms, for $t \in \mathbb{Z}_{\geq 0}$,

$$(2.8) \quad p(t+1) = \text{Cd}(V(p(t))).$$

A variation of the function $\mathcal{H}_{\text{multicenter}}$ is useful to analyze this algorithm. We define the positions-based multicenter function $\mathcal{H}_{\text{Voronoi}} : Q^N \setminus S_N \rightarrow \mathbb{R}_{\geq 0}$ by

$$(2.9) \quad \mathcal{H}_{\text{Voronoi}}(p) = \mathcal{H}_{\text{multicenter}}(V(p), p) = \sum_{i=1}^N \int_{V_i(p)} f(\|q - p_i\|) \phi(q) dq.$$

Because of the compactness of Q , a continuity property, and the monotonicity properties in Proposition 2.4, one can show [10, Theorem 5.5] that $\mathcal{H}_{\text{Voronoi}}$ is monotonically non-increasing along the solutions of (2.8) and that all solutions of (2.8) converge asymptotically to the set of configurations that generate centroidal Voronoi partitions. Additional considerations about convergence are given in [14].

3. Gossip coverage control as a dynamical system on the space of partitions. In this section we present the problem of interest, our novel gossip coverage algorithm and its convergence properties in Subsections 3.1, 3.2 and 3.3, respectively. In order to reduce the communication requirements of our algorithm, we propose an adjacency-based and continuous algorithm in Subsection 3.4. Finally, we report some simulation results in Subsection 3.5.

3.1. Problem statement. The distributed coverage law, based upon the Lloyd algorithm and described in the previous section, has some important limitations: it is applicable only to robotic networks with *synchronized* and *reliable* communication along *all edges of the Delaunay graph* (computed as a function of the robots’ positions). In other words, the law (2.8) requires that there exists a predetermined common communication schedule for all robots and, at each communication round, each robot must simultaneously and reliably communicate its position. Note that the Delaunay graph, interpreted as a communication graph, has the following drawbacks: for worst-case robots’ positions, a robot might have $N - 1$ neighbors in the Delaunay graph and/or might have a neighbor that is arbitrarily far inside the environment. Therefore,

each robot must be capable to communicate potentially to all other robots and/or to robots at large distances.

Given this broad range of undesirable limitations, the aim of this paper is to reduce the communication requirements of distributed coverage algorithms, in terms of reliability, synchronization and topology. Here are some relevant questions that constitute our informal problem statement:

Is it possible to optimize robots positions and environment partition with asynchronous, unreliable, and delayed communication? Specifically, what if the communication model is that of *gossiping agents*, that is, a model in which only a pair of robots can communicate at any time? Since Voronoi partitions generated by gossiping and moving agents cannot be computed by gossiping agents, how do we update the environment partition?

To answer these questions, the next subsections propose an innovative partition-based gossip approach, in which the robots' positions essentially play no role and where instead dominance regions are iteratively updated. Designing coverage algorithms as dynamical systems on the space of partitions has the key advantage that one is not restricted to working only with Voronoi or anyway position-dependent partitions.

EXAMPLE 3.1 (The Lloyd algorithm in the partition-based approach). *The distributed coverage algorithm (2.8) updates the robots' positions so as to incrementally minimize the function $\mathcal{H}_{\text{Voronoi}}$, while the environment partition is a function of the robots' positions. In this paper we take a dual approach: we consider an algorithm that evolves partitions. From this partition-based viewpoint, the coverage algorithm is an iterated map on \mathcal{V}_N and equation (2.8) is rewritten as $v(t+1) = V(\text{Cd}(v(t)))$. \square*

3.2. The gossip coverage algorithm. In this subsection we present a novel partition-based coverage algorithm in which, at each communication round, only two regions communicate. Recall the notion of bisector half-space from equation (2.3).

Gossip Coverage Algorithm

For all $t \in \mathbb{Z}_{\geq 0}$, each agent $i \in \{1, \dots, N\}$ maintains in memory a dominance region $v_i(t)$. The collection $(v_1(0), \dots, v_N(0))$ is an arbitrary polygonal N -partition of Q . At each $t \in \mathbb{Z}_{\geq 0}$ a pair of communicating regions, say $v_i(t)$ and $v_j(t)$, is selected by a deterministic or stochastic process to be determined. Every agent $k \notin \{i, j\}$ sets $v_k(t+1) := v_k(t)$. Agents i and j perform the following tasks:

- 1: agent i transmits to agent j its dominance region $v_i(t)$ and vice-versa
 - 2: both agents compute the centroids $\text{Cd}(v_i(t))$ and $\text{Cd}(v_j(t))$
 - 3: **if** $\text{Cd}(v_i(t)) = \text{Cd}(v_j(t))$ **then**
 - 4: $v_i(t+1) := v_i(t)$ and $v_j(t+1) := v_j(t)$
 - 5: **else**
 - 6: $v_i(t+1) := (v_i(t) \cup v_j(t)) \cap H_{\text{bisector}}(\text{Cd}(v_i(t)); \text{Cd}(v_j(t)))$
 $v_j(t+1) := (v_i(t) \cup v_j(t)) \cap H_{\text{bisector}}(\text{Cd}(v_j(t)); \text{Cd}(v_i(t)))$
-

In other words, when two agents with distinct centroids communicate, their dominance regions evolve as follows: the union of the two dominance regions is divided into two new dominance regions by the hyperplane bisecting the segment between the two centroids; see Fig. 3.1. As a consequence, if the centroids $\text{Cd}(v_i(t))$, $\text{Cd}(v_j(t))$ are distinct, then $\{v_i(t+1), v_j(t+1)\}$ is the Voronoi partition of the set $v_i(t) \cup v_j(t)$ generated by the centroids $\text{Cd}(v_i(t))$ and $\text{Cd}(v_j(t))$.

We claim that the algorithm is well-posed in the sense that the sequence of collections $\{v(t)\}_{t \in \mathbb{Z}_{\geq 0}}$ generated by the algorithm is an N -partition at all times t . Indeed,




FIG. 3.1. The gossip coverage algorithm. The left and right figure contain the initial partition and the partition after one application of the gossip coverage algorithm. In the middle figure we show the two centroids and (with a dashed line) the segment determining the bisector half-space.

it is immediate to see that the first two properties in Definition 2.1 are satisfied at all time if they are satisfied at initial time. Finally, at all times t , each component of $v(t)$ is clearly closed and has non-empty interior because of the following geometric fact: there exists no half-plane containing the interior of a region and not containing the centroid of the same region.

Now, for any $i, j \in \{1, \dots, N\}$ with $i \neq j$, define the map $T_{ij} : \mathcal{V}_N \rightarrow \mathcal{V}_N$ by

$$T_{ij}(v) = \begin{cases} v, & \text{if } \text{Cd}(v_i) = \text{Cd}(v_j), \\ (v_1, \dots, \widehat{v}_i, \dots, \widehat{v}_j, \dots, v_N), & \text{otherwise,} \end{cases}$$

where

$$(3.1) \quad \begin{aligned} \widehat{v}_i &= (v_i \cup v_j) \cap H_{\text{bisector}}(\text{Cd}(v_i); \text{Cd}(v_j)), \\ \widehat{v}_j &= (v_i \cup v_j) \cap H_{\text{bisector}}(\text{Cd}(v_j); \text{Cd}(v_i)). \end{aligned}$$

The dynamical system on the space of partitions is therefore described by, for $t \in \mathbb{Z}_{\geq 0}$,

$$(3.2) \quad v(t+1) = T_{ij}(v(t)),$$

together with a rule describing what pair of regions (i, j) is selected at each time. We also define the set-valued map $T : \mathcal{V}_N \rightrightarrows \mathcal{V}_N$ by $T(v) = \{T_{ij}(v) \mid i, j \in \{1, \dots, N\}, i \neq j\}$. The map T describes one iteration of the gossip coverage algorithm; an evolution of the gossip coverage algorithm is one of the solutions to the non-deterministic set-valued dynamical system $v(t+1) \in T(v(t))$.

REMARK 3.2 (Gossip disk-covering control). *We believe that our gossip and partition-based algorithmic approach is applicable to a broad range of coverage problems. For example, the worst-case multicenter function [10] is defined by $\mathcal{H}_{\text{worst}}(p, v) = \max_{i \in \{1, \dots, N\}} \max_{q \in v_i} \|q - p\|$. Maximizing $\mathcal{H}_{\text{worst}}$ is equivalent to covering Q with N disks of smallest radius centered at p . As in Proposition 2.4, for any $v \in \mathcal{V}_N$ and $p \in Q^N \setminus S_N$, one can prove $\mathcal{H}_{\text{worst}}(V(p), p) \leq \mathcal{H}_{\text{worst}}(v, p)$ and $\mathcal{H}_{\text{worst}}(v, \text{CC}(v)) \leq \mathcal{H}_{\text{worst}}(v, p)$, where $\text{CC}(v)$ is the array of circumcenters of the components of v . Hence, a gossip coverage algorithm for $\mathcal{H}_{\text{worst}}$ is designed by replacing centroid with circumcenter operations in (3.1). We leave this and further extensions to future works. \square*

3.3. Analysis results for the gossip coverage algorithm. In this subsection we state the main analysis and convergence results for the gossip coverage algorithm.

We begin by studying the fixed points of T and by introducing an appropriate cost function with monotonicity properties along T . Regarding the algorithm's fixed

points, we extend Definition 2.5 as follows. A partition v is *mixed centroidal Voronoi* if, for all pairs (v_i, v_j) with $i \neq j$, either $\text{Cd}(v_i) = \text{Cd}(v_j)$ or (v_i, v_j) is a centroidal Voronoi partition of $v_i \cup v_j$, that is, $v_i = (v_i \cup v_j) \cap H_{\text{bisector}}(\text{Cd}(v_i); \text{Cd}(v_j))$.

LEMMA 3.3 (Fixed points of T and centroidal Voronoi partitions). *For $i, j \in \{1, \dots, N\}$, $j \neq i$, denote the set of fixed points of $T_{ij} : \mathcal{V}_N \rightarrow \mathcal{V}_N$ by $F_{ij} = \{v \in \mathcal{V}_N \mid T_{ij}(v) = v\}$. The following statements hold:*

- (i) $\bigcap_{j \neq i} F_{ij}$ equals the set of mixed centroidal Voronoi partitions; and
- (ii) if v is a mixed centroidal Voronoi partition satisfying $\text{Cd}(v_i) \neq \text{Cd}(v_j)$ for $j \neq i$, then v is centroidal Voronoi.

Next, we define the partition-based multicenter function $\mathcal{H}_{\text{centroid}} : \mathcal{V}_N \rightarrow \mathbb{R}_{\geq 0}$ by

$$(3.3) \quad \mathcal{H}_{\text{centroid}}(v) = \mathcal{H}_{\text{multicenter}}(v, \text{Cd}(v)) = \sum_{i=1}^N \int_{v_i} f(\|q - \text{Cd}(v_i)\|) \phi(q) dq.$$

LEMMA 3.4 (Monotonicity of $\mathcal{H}_{\text{centroid}}$ along T). *For $i, j \in \{1, \dots, N\}$, $i \neq j$,*

$$\begin{aligned} \mathcal{H}_{\text{centroid}}(T_{ij}(v)) &\leq \mathcal{H}_{\text{centroid}}(v), & \text{for all } v \in \mathcal{V}_N, \text{ and} \\ \mathcal{H}_{\text{centroid}}(T_{ij}(v)) &< \mathcal{H}_{\text{centroid}}(v), & \text{iff } T_{ij}(v) \text{ and } v \text{ differ by a set of measure zero.} \end{aligned}$$

The proofs of Lemmas 3.3 and 3.4 consist of elementary manipulations and are omitted in the interest of brevity. In short, we have established that the function $\mathcal{H}_{\text{centroid}}$ monotonically decreases along each T_{ij} when away from fixed points, and that centroidal Voronoi partitions are the fixed points of all T_{ij} provided centroids are distinct.

We now prepare to state the main convergence result for T . We need to introduce some useful properties for sequences of partitions and for switching signals.

DEFINITION 3.5 (Non-degeneracy). *A sequence of N -partitions $\{v(t)\}_{t \in \mathbb{Z}_{\geq 0}}$ is*

- (i) (uniformly) distinct centroidal if there exists $\epsilon > 0$ such that, for all $t \in \mathbb{Z}_{\geq 0}$ and $i, j \in \{1, \dots, N\}$, $i \neq j$, one has $\|\text{Cd}(v_i(t)) - \text{Cd}(v_j(t))\| \geq \epsilon$;
- (ii) (uniformly componentwise) non-vanishing if there exists $\epsilon > 0$ such that, for all $t \in \mathbb{Z}_{\geq 0}$ and $i \in \{1, \dots, N\}$, the Lebesgue measure of $v_i(t)$ is greater than ϵ ; and
- (iii) (uniformly componentwise) finitely convex if there exists $\ell \in \mathbb{N}$ such that, for all $t \in \mathbb{Z}_{\geq 0}$ and $i \in \{1, \dots, N\}$, the set $v_i(t)$ is the union of at most ℓ convex sets.

Moreover, the sequence v is said to be non-degenerate if it is distinct centroidal, non-vanishing and finitely convex.

For example, a sequence of partitions is finitely-convex if each component of each partition in the sequence is the union of a uniformly bounded number of polygons with a uniformly bounded number of vertices.

DEFINITION 3.6 (Uniform and random persistency). *Let X be a finite set.*

- (i) A map $\sigma : \mathbb{Z}_{\geq 0} \rightarrow X$ is uniformly persistent if there exists a duration $\Delta \in \mathbb{N}$ such that, for each $x \in X$, there exists an increasing sequence of times $\{t_k\}_{k \in \mathbb{Z}_{\geq 0}} \subset \mathbb{Z}_{\geq 0}$ satisfying $t_{k+1} - t_k \leq \Delta$ and $\sigma(t_k) = x$ for all $k \in \mathbb{Z}_{\geq 0}$.
- (ii) A stochastic process $\sigma : \mathbb{Z}_{\geq 0} \rightarrow X$ is randomly persistent if there exists a probability $p \in]0, 1[$ and a duration $\Delta \in \mathbb{N}$ such that, for each $x \in X$ and for each $t \in \mathbb{Z}_{\geq 0}$, there exists $k \in \{1, \dots, \Delta\}$ satisfying

$$\mathbb{P}[\sigma(t+k) = x \mid \sigma(t), \dots, \sigma(1)] \geq p.$$

We are now ready to state the main deterministic and stochastic convergence results for the gossip coverage algorithm. It is convenient to postpone to Section 6.2 the theorem proof and the definition of convergence in the space of partitions.

THEOREM 3.7 (Convergence under persistent gossip). *Consider the gossip coverage algorithm T and the evolutions $\{v(t)\}_{t \in \mathbb{Z}_{\geq 0}} \subset \mathcal{V}_N$ defined by*

$$v(t+1) = T_{\sigma(t)}(v(t)), \quad \text{for } t \in \mathbb{Z}_{\geq 0},$$

where $\sigma : \mathbb{Z}_{\geq 0} \rightarrow \{(i, j) \in \{1, \dots, N\}^2 \mid i \neq j\}$ is either a deterministic map or a stochastic process. Then the following statements hold:

- (i) if σ is a uniformly persistent map, then each non-degenerate evolution v converges to the set of centroidal Voronoi partitions; and
- (ii) if σ is a randomly persistent stochastic process, then each evolution v , conditioned upon being non-degenerate, converges almost surely to the set of centroidal Voronoi partitions.

We conjecture that evolutions resulting from initial polygonal partitions are always non-degenerate. Section 3.5 contains some numerical evidence to this effect.

Lemma 3.4 indicates how the function $\mathcal{H}_{\text{centroid}}$ plays the role of a Lyapunov function for the dynamical system defined by T . To provide a complete Lyapunov convergence proof of Theorem 3.7, we set out to establish three sets of relevant results. First, we need to establish extensions of the Krasovskii-LaSalle invariance principle for set-valued dynamical systems over compact metric spaces. Second, we need to establish the compactness properties of the space of non-degenerate partitions. Third, we need to establish the continuity of the relevant geometric maps. These three topics are the subjects of Section 4, 5 and 6, respectively.

3.4. Designing an adjacency-based and continuous algorithm. The gossip coverage map T has the undesirable feature that it entails communication exchanges between any two regions. This communication requirement might be too onerous in some multiagent applications. Ideally we would like to require communications only between *adjacent* regions, i.e., regions whose boundaries touch, or between “nearby” regions. We believe such communication requirements may be easily achieved in practice; for completeness purposes, a sample implementation for robots with range-dependent unreliable communication is presented in the report [9]. Hence, we wish to design a modification of the map T which acts only on adjacent regions, taking into account that the modified update map needs to be continuous for technical reasons: the invariance principles we adopt for the convergence analysis require continuity of the dynamical system.

Motivated by this discussion, we modify the map T to rely upon only adjacency-based communication and to be continuous. First, we introduce a pseudodistance notion between sets. Given closed $A \subset Q$ and $B \subset Q$ with non-empty interior, define

$$\text{pseudodist}(A, B) = \inf\{\|a - b\| \mid (a, b) \in \text{int}(A) \times \text{int}(B)\}.$$

Second, we select a positive constant $\delta \ll \text{diam}(Q)$ and denote by $T^\delta : \mathcal{V}_N \rightarrow \mathcal{V}_N$ the *modified gossip coverage map* to be defined in what follows. For any $i, j \in \{1, \dots, N\}$, $i \neq j$, we give the following partial definition:

$$(3.4) \quad T_{ij}^\delta(v) = \begin{cases} v, & \text{if } (\| \text{Cd}(v_i) - \text{Cd}(v_j) \| = 0) \text{ or } (\text{pseudodist}(v_i, v_j) \geq \delta), \\ T(v), & \text{if } (\| \text{Cd}(v_i) - \text{Cd}(v_j) \| \geq \delta) \text{ and } (\text{pseudodist}(v_i, v_j) = 0). \end{cases}$$

Therefore, if either $\text{Cd}(v_i)$ and $\text{Cd}(v_j)$ coincide or the pseudodistance between v_i and v_j is larger than δ , then $T_{ij}^\delta(v) = v$, that is, the map T_{ij}^δ leaves the partition unchanged. Additionally, if the pseudodistance between the regions v_i and v_j is zero (v_i and v_j are adjacent) and the distance between $\text{Cd}(v_i)$ and $\text{Cd}(v_j)$ is larger than δ , then $T_{ij}^\delta(v) = T_{ij}(v)$.

Next, we consider partitions that do not satisfy either of the two conditions in definition (3.4). We define the *unit saturation function* $\text{sat} : \mathbb{R}_{\geq 0} \rightarrow [0, 1]$ by $\text{sat}(x) = x$ if $x \in [0, 1]$, and $\text{sat}(x) = 1$ if $x > 1$ and the scaling function $\beta_{ij} : \mathcal{V}_N \rightarrow [0, 1]$ by

$$\beta_{ij}(v) = \text{sat}(\|\text{Cd}(v_i) - \text{Cd}(v_j)\|/\delta) \left(1 - \text{sat}(\text{pseudodist}(v_i, v_j)/\delta)\right).$$

The first condition and the second condition in (3.4) correspond precisely to $\beta_{ij}(v) = 0$ and $\beta_{ij}(v) = 1$, respectively. For partitions v satisfying $0 < \beta_{ij}(v) < 1$, we aim to define T^δ so as to continuously interpolate between the identity map and the map T ; see Fig. 3.2 for an illustration. Let $R_i = v_i \cap H_{\text{bisector}}(\text{Cd}(v_j), \text{Cd}(v_i))$ and




FIG. 3.2. *Modified gossip between close but not adjacent regions ($0 < \beta_{12}(v) < 1$). The bisecting line γ_\perp borders the set $R_2 = v_2 \cap H_{\text{bisector}}(\text{Cd}(v_1), \text{Cd}(v_2))$ that, in the map T , is assigned to v_1 (see Fig. 3.1). According to T^δ instead, only the subset $\widehat{R}_2 \subsetneq R_2$ is assigned to v_1 . Loosely speaking, the “width” of \widehat{R}_2 equals $\beta_{12}(v)$ times the “width” of R_2 , where “width” of a set is the maximum distance from a point in the set to γ_\perp .*

$R_j = v_j \cap H_{\text{bisector}}(\text{Cd}(v_i), \text{Cd}(v_j))$. Define the line $\gamma_\perp = \partial H_{\text{bisector}}(\text{Cd}(v_j), \text{Cd}(v_i))$ and

$$(3.5) \quad \begin{aligned} \widehat{p}_i &= \text{a point in } \overline{(R_i)} \text{ that is maximally distant from } \gamma_\perp, \\ \widehat{p}_j &= \text{a point in } \overline{(R_j)} \text{ that is maximally distant from } \gamma_\perp. \end{aligned}$$

Next, note that for each $q \in R_i \cup R_j$ there exists a unique line, say γ_q , that is parallel to γ_\perp and passes through q . Based on this notion, we define

$$\begin{aligned} \widehat{R}_i &= \{q \in R_i \mid \text{dist}(\widehat{p}_i, \gamma_q) \leq \beta_{ij}(v) \text{dist}(\widehat{p}_i, \gamma_\perp) \text{ or } \text{dist}(q, \gamma_\perp) \geq \text{dist}(\widehat{p}_i, \gamma_\perp)\}, \\ \widehat{R}_j &= \{q \in R_j \mid \text{dist}(\widehat{p}_j, \gamma_q) \leq \beta_{ij}(v) \text{dist}(\widehat{p}_j, \gamma_\perp) \text{ or } \text{dist}(q, \gamma_\perp) \geq \text{dist}(\widehat{p}_j, \gamma_\perp)\}. \end{aligned}$$

We can now complete the partial definition (3.4). For all v with $0 < \beta_{ij}(v) < 1$, that is, for all partitions not already dealt with in definition (3.4), we define

$$T_{ij}^\delta(v) = (v_1, \dots, \underbrace{(v_i \setminus \widehat{R}_i) \cup \widehat{R}_i}_{\text{ith entry}}, \dots, \underbrace{(v_j \setminus \widehat{R}_j) \cup \widehat{R}_j}_{\text{jth entry}}, \dots, v_N).$$

As discussed for T , one can prove that the map $T^\delta : \mathcal{V}_N \rightrightarrows \mathcal{V}_N$ defined by $T^\delta(v) = \{T_{ij}^\delta(v) \mid i, j \in \{1, \dots, N\}, i \neq j\}$, is well-posed and has the following properties.

THEOREM 3.8 (Convergence of modified gossip map). *Consider the modified gossip coverage algorithm T^δ and the evolutions $\{v(t)\}_{t \in \mathbb{Z}_{\geq 0}} \subset \mathcal{V}_N$ defined by*

$$v(t+1) = T_{\sigma(t)}^\delta(v(t)), \quad \text{for } t \in \mathbb{Z}_{\geq 0},$$

where $\sigma : \mathbb{Z}_{\geq 0} \rightarrow \{(i, j) \in \{1, \dots, N\}^2 \mid i \neq j\}$ is either a deterministic map or a stochastic process. Then the following statements hold:

- (i) if σ is a uniformly persistent map, then each non-vanishing and finitely-convex evolution v converges to the set of mixed centroidal Voronoi partitions; and
- (ii) if σ is a randomly persistent stochastic process, then each evolution v , conditioned upon being non-vanishing and finitely convex, converges almost surely to the set of mixed centroidal Voronoi partitions.

3.5. Simulation results and implementation remarks. We have extensively simulated the partition-based gossip coverage algorithm T on a 2-dimensional polygonal environment with uniform density and performance function $f(x) = x^2$. Simulations have been implemented as a `Matlab` program, using the `General Polygon Clipper Library` to perform operations on polygons. We adopted the following communication model: at each iteration, a region pair is chosen, uniformly at random, among all pairs of adjacent regions. Fig. 3.3 is an illustration of a typical evolution.




FIG. 3.3. An example simulation of the gossip coverage algorithm with uniform random edge selection. The environment Q is a rectangle with uniform density, centroids are computed with performance function $f(x) = x^2$, and $N = 6$ regions compose the partition. Snapshots of an evolution are shown for $t \in \{0, 20, 50, 100, 300\}$. One may verify numerically that the sequence converges asymptotically to a centroidal Voronoi partition. At $t = 20$ one of the regions is disconnected.

Our first numerical finding is that the gossip coverage algorithm appears to converge to centroidal Voronoi partitions from all initial conditions. This is the same property that the Lloyd synchronous coverage algorithm (2.8) is known to possess. In other words, our numerically-computed sequences of partitions always converge to centroidal Voronoi partitions – even though our theoretical analysis (1) requires a continuous interpolation from T to T^δ and (2) does not exclude convergence to degenerate partitions where some component regions might have coincident centroids, or might have empty interiors, or might be composed of “polygons with an infinite number of vertices,” that is, arbitrary sets.

A second numerical finding is that, throughout numerous sample executions, the polygonal regions, which appear during the evolution, rarely have complicated shapes and large numbers of vertices. This can be intuitively explained by the remark that whenever two adjacent agents communicate, the border between their regions becomes a straight line. This limited complexity is good news because of our assumption of finite convexity and because large numbers of vertices affect both the computation and the communication burden of the gossip coverage algorithm.

Finally, it is possible, and we have observed it numerically, to have evolutions of the algorithm that, before converging to centroidal Voronoi partitions, have components with disconnected regions. We believe that there might be applications where it

is desirable to maintain connectivity of the components of the partition and, therefore, we sketch here how to design a connectivity-preserving algorithm. Note that the update step of the partition-based coverage algorithm amounts to the exchange, among the agents, of a region, which consists in general of several connected components. In the connectivity-preserving algorithm, such components are considered individually, and each of them is traded only if this can be done without losing connectivity; if not, the component is kept by the previous owner. Numerical simulations indicate that such an algorithm leads to centroidal Voronoi partitions as well.

4. On the Krasovskii–LaSalle invariance principle: set-valued maps on metric spaces. In this section we consider discrete-time set-valued dynamical system defined on metric spaces. Our goal is to provide some extensions of the classical Krasovskii–LaSalle Invariance Principle; we refer the interested reader to [5, 22, 35] for recent invariance principles for switched continuous-time and hybrid systems.

We start by reviewing some preliminary notions including set-valued dynamical systems, continuity and invariance properties, and Lyapunov functions. On a metric space (X, d) , where X is a set and d is a metric on X , a set-valued map $T : X \rightrightarrows X$ is non-empty if $T(x) \neq \emptyset$ for all $x \in X$. An evolution of the dynamical system determined by a non-empty set-valued map T is a sequence $\{x_n\}_{n \in \mathbb{Z}_{\geq 0}} \subset X$ with the property that

$$x_{n+1} \in T(x_n), \quad n \in \mathbb{Z}_{\geq 0}.$$

In other words, we regard a set-valued map as a nondeterministic discrete-time dynamical system. For set-valued maps we introduce notions of continuity and invariance as follows. A set-valued map T is *closed* at $x \in X$ if, for all pairs of convergent sequences $x_k \rightarrow x$ and $x'_k \rightarrow x'$ such that $x'_k \in T(x_k)$, one has that $x' \in T(x)$. Additionally, T is closed on $W \subset X$ if it is closed at all $w \in W$. A set $W \subset X$ is *weakly positively invariant* for T if $T(w) \cap W$ is non-empty for all $w \in W$. A set W is *strongly positively invariant* for T if $T(w) \subset W$ for all $w \in W$.

We are ready now to state a Krasovskii–LaSalle invariance principle for set-valued maps defined on metric spaces. This result extends the Global Convergence Theorem in [27] to more general Lyapunov functions. Its proof follows the lines of the proof of Theorem 1.21 in [10], and is thus omitted.

LEMMA 4.1 (Krasovskii–LaSalle invariance principle for set-valued maps). *Let (X, d) be a metric space and $T : X \rightrightarrows X$ be non-empty. Assume that:*

- (i) *there exists a compact set $W \subseteq X$ that is strongly positively invariant for T ;*
- (ii) *there exists a function $U : W \rightarrow \mathbb{R}$ such that $U(w') \leq U(w)$, for all $w \in W$ and $w' \in T(w)$;*
- (iii) *the function U is continuous on W and the map T is closed on W .*

Then there exists $c \in \mathbb{R}$ such that each evolution of T with initial condition in W approaches a set of the form $S \cap U^{-1}(c)$, where S is the largest weakly positively invariant set contained in

$$\{w \in W \mid \exists w' \in T(w) \text{ such that } U(w') = U(w)\}.$$

In this paper, given the metric space (X, d) , we deal with a set-valued map $T : X \rightrightarrows X$ defined by a collection of maps $T_1, \dots, T_m : X \rightarrow X$ via the equality $T(x) = \{T_1(x), \dots, T_m(x)\}$ for $x \in X$. For this kind of set-valued maps, closedness is related to continuity of ordinary maps.

LEMMA 4.2. *Let $T_1, \dots, T_m : X \rightarrow X$ be continuous on $W \subset X$. The set-valued map $T : X \rightrightarrows X$ defined by $T(x) = \{T_1(x), \dots, T_m(x)\}$ is closed on W .*

Proof. Let $w_n \rightarrow w$ and $w'_n \rightarrow w'$ be a pair of convergent sequences in W , such that $w'_n \in T(w_n)$. We claim that the continuity of T_1, \dots, T_m implies $w' \in T(w)$.

Note that, by hypothesis, for all $n \in \mathbb{Z}_{\geq 0}$ there exists $i_n \in \{1, \dots, m\}$ such that $w'_n = T_{i_n}(w_n)$. Because the set $\{1, \dots, m\}$ is finite, there exists an index $j \in \{1, \dots, m\}$ that appears infinitely many times in the sequence $\{i_n\}_{n \in \mathbb{N}}$. Consider the subsequences $\{w_{n_i}\} \subseteq \{w_n\}$ and $\{w'_{n_i}\} \subseteq \{w'_n\}$, such that $w'_{n_i} = T_j(w_{n_i})$. Clearly, we have that $w_{n_i} \rightarrow w$ and $w'_{n_i} \rightarrow w'$, where from the continuity of T_j it follows that $w' = T_j(w)$. Thus, $w' \in T(w)$ and the claim is proved. \square

The following result is a stronger version of Lemma 4.1, for a particular class of set-valued dynamical systems.

THEOREM 4.3 (Uniformly persistent switches imply convergence). *Let (X, d) be a metric space. Given a collection of maps $T_1, \dots, T_m : X \rightarrow X$, define the set-valued map $T : X \rightrightarrows X$ by $T(x) = \{T_1(x), \dots, T_m(x)\}$ and let $\{x_n\}_{n \in \mathbb{Z}_{\geq 0}}$ be an evolution of T . Assume that:*

- (i) *there exists a compact set $W \subseteq X$ that is strongly positively invariant for T ;*
- (ii) *there exists a function $U : W \rightarrow \mathbb{R}$ such that $U(w') < U(w)$, for all $w \in W$ and $w' \in T(w) \setminus \{w\}$;*
- (iii) *the maps T_i , for $i \in \{1, \dots, m\}$, and U are continuous on W ; and*
- (iv) *for all $i \in \{1, \dots, m\}$, there exists an increasing sequence of times $\{n_k \mid k \in \mathbb{Z}_{\geq 0}\}$ such that $x_{n_{k+1}} = T_i(x_{n_k})$ and $(n_{k+1} - n_k)$ is bounded.*

If $x_0 \in W$, there exists $c \in \mathbb{R}$ such that the evolution $\{x_n\}_{n \in \mathbb{Z}_{\geq 0}}$ approaches the set

$$(F_1 \cap \dots \cap F_m) \cap U^{-1}(c),$$

where $F_i = \{w \in W \mid T_i(w) = w\}$ is the set of fixed points of T_i in W , $i \in \{1, \dots, m\}$.

Loosely speaking, (i) the compactness of a strongly positively invariant set, (ii) a monotonicity property for a Lyapunov function, (iii) continuity properties, and (iv) uniformly persistent switches among finitely many maps, together ensure convergence of each evolution to the intersection of the fixed points of the maps.

Proof. [Proof of Theorem 4.3] Let S be the largest weakly positively invariant set contained in

$$\{w \in W \mid \exists w' \in T(w) \text{ such that } U(w') = U(w)\} = F_1 \cup \dots \cup F_m.$$

Since T is closed by Lemma 4.2, the assumptions of Lemma 4.1 are met; hence there exists $c \in \mathbb{R}$ such that $U(x_n) \rightarrow c$ and $x_n \rightarrow S \cap U^{-1}(c)$.

Let $\omega(x_n)$ denote the ω -limit set of the sequence $\{x_n \mid n \in \mathbb{Z}_{\geq 0}\}$. If we show that $\omega(x_n) \subseteq (F_1 \cap \dots \cap F_m) \cap U^{-1}(c)$, then the statement of the theorem is proved. We proceed by contradiction. To this aim, let $\hat{x} \in S \cap U^{-1}(c) \setminus ((F_1 \cap \dots \cap F_m) \cap U^{-1}(c))$ and let $\{x_{n_h}\}_{h \in \mathbb{Z}_{\geq 0}}$ be a subsequence such that $x_{n_h} \rightarrow \hat{x}$.

Observe that for each $\hat{x} \in S \setminus (F_1 \cap \dots \cap F_m)$, there exists a non-empty set $\mathcal{I}_{\hat{x}} \subset \{1, \dots, m\}$ such that, $\hat{x} = T_i(\hat{x})$ if $i \in \mathcal{I}_{\hat{x}}$, and $\hat{x} \neq T_i(\hat{x})$ if $i \notin \mathcal{I}_{\hat{x}}$. By the continuity of the maps T_i , there exists $\delta \in \mathbb{R}_{>0}$ such that, if $i \notin \mathcal{I}_{\hat{x}}$, then $T_i(x) \neq x$ for all $x \in B_\delta(\hat{x}) = \{x \in W \mid d(x, \hat{x}) \leq \delta\}$. Let now

$$\gamma_\delta = \min_{i \in \mathcal{I}_{\hat{x}}} \left\{ \min_{x \in B_\delta(\hat{x})} \left(U(x) - U(T_i(x)) \right) \right\} \geq 0.$$

By hypothesis, if $i \notin \mathcal{I}_{\hat{x}}$, then $U(T_i(x)) < U(x)$ for all $x \in B_\delta(\hat{x})$. Hence, since $B_\delta(\hat{x})$ is closed, and U and the maps T_i are continuous, we deduce that $\gamma > 0$.

Observe now that hypothesis (iv) implies the existence of a duration $D \in \mathbb{N}$ such that every map T_i , $i \in \{1, \dots, m\}$, is applied at least once within every interval $[n, n + D[$, for all $n \in \mathbb{Z}_{\geq 0}$. Consider the set $\{T_i\}_{i \in \mathcal{I}_{\hat{x}}}$; this is a collection of continuous maps having \hat{x} as fixed point. Then, there exists a suitable $\epsilon \in \mathbb{R}_{>0}$ such that, given any r -uple $(j_1, \dots, j_r) \in \mathcal{I}_{\hat{x}}^r$, $r \leq D$, we have that $T_{j_1} \circ T_{j_2} \circ T_{j_3} \circ \dots \circ T_{j_r}(x) \in B_\delta(\hat{x})$ for all $w \in B_\epsilon(\hat{x})$.

Select now k such that the element x_{n_k} in the subsequence $\{x_{n_h}\}_{h \in \mathbb{Z}_{\geq 0}}$ satisfies $d(x_{n_k}, \hat{x}) < \epsilon$ and $U(x_{n_k}) - c < \gamma_\delta$. Let

$$s = \min\{t \in [1, D[\mid \exists j \notin \mathcal{I}_{\hat{x}} \text{ such that } x_{n_k+t+1} = T_j(x_{n_k+t})\}.$$

Observe that $U(x_{n_k+s}) - c < \gamma_\delta$ and $U(x_{n_k+s}) - U(T_j(x_{n_k+s})) \geq \gamma_\delta$ implying that $U(T_j(x_{n_k+s})) < c$. This is a contradiction. \square

REMARK 4.4. *An alternate proof of this theorem can be given by applying an invariance principle obtained in [35] on an appropriately-designed dynamical extension of the discrete-time set-valued system.* \square

In Appendix A we show how the persistent switching assumption is necessary. Next, we provide a probabilistic version of the previous theorem.

THEOREM 4.5 (Persistent random switches imply convergence). *Let (X, d) be a metric space. Given a collection of maps $T_1, \dots, T_m : X \rightarrow X$, define the set-valued map $T : X \rightrightarrows X$ by $T(x) = \{T_1(x), \dots, T_m(x)\}$. Given a stochastic process $\sigma : \mathbb{Z}_{\geq 0} \rightarrow \{1, \dots, m\}$, consider an evolution $\{x_n\}_{n \in \mathbb{Z}_{\geq 0}}$ of T satisfying*

$$x_{n+1} = T_{\sigma(n)}(x_n).$$

Assume that:

- (i) *there exists a compact set $W \subseteq X$ that is strongly positively invariant for T ;*
- (ii) *there exists a function $U : W \rightarrow \mathbb{R}$ such that $U(w') < U(w)$, for all $w \in W$ and $w' \in T(w) \setminus \{w\}$;*
- (iii) *the maps T_i , for $i \in \{1, \dots, m\}$, and U are continuous on W ; and*
- (iv) *there exists $p \in]0, 1[$ and $k \in \mathbb{N}$ such that, for all $i \in \{1, \dots, m\}$ and $n \in \mathbb{Z}_{\geq 0}$, there exists $h \in \{1, \dots, k\}$ such that*

$$\mathbb{P}[\sigma(n+h) = i \mid \sigma(n), \dots, \sigma(1)] \geq p.$$

If $x_0 \in W$, then there exists $c \in \mathbb{R}$ such that almost surely the evolution $\{x_n\}_{n \in \mathbb{Z}_{\geq 0}}$ approaches the set

$$(F_1 \cap \dots \cap F_m) \cap U^{-1}(c),$$

where $F_i = \{w \in W \mid T_i(w) = w\}$ is the set of fixed points of T_i in W , $i \in \{1, \dots, m\}$.

Loosely speaking, (i) the compactness of a strongly positively invariant set, (ii) a monotonicity property for a Lyapunov function, (iii) continuity properties, and (iv) persistent random switches among finitely many maps, together ensure convergence of each evolution to the intersection of the fixed points of the maps.

Proof. [Proof of Theorem 4.5] If $x_0 \in W$, then the stochastic process σ induces a stochastic process taking values in W . From now on, we restrict our attention to sequences $\{x_n\}_{n \in \mathbb{Z}_{\geq 0}}$ such that $x_0 \in W$. In other words we assume that the sample space containing all the evolutions of our interest is given by

$$\mathcal{A} = \{\{x_n\}_{n \in \mathbb{Z}_{\geq 0}} \mid x_n \in W \text{ for all } n \in \mathbb{Z}_{\geq 0}\}.$$

Let S be the largest weakly positively invariant set contained in

$$\{w \in W \mid \exists w' \in T(w) \text{ such that } U(w') = U(w)\} = F_1 \cup \dots \cup F_m.$$

From Lemma 4.1, we know that there exists $c \in \mathbb{R}$ such that $x_n \rightarrow S \cap U^{-1}(c)$. This implies that the following event is certain:

$$E = \{\{x_n\}_{n \in \mathbb{Z}_{\geq 0}} \mid \exists c \in \mathbb{R} \text{ such that } \lim_{n \rightarrow \infty} U(x_n) = c\}.$$

Let $\omega(x_n)$ denote the ω -limit set of the sequence $\{x_n \mid n \in \mathbb{Z}_{\geq 0}\}$. If we show that $\omega(x_n) \subseteq ((F_1 \cap \dots \cap F_m) \cap U^{-1}(c))$ almost surely, then the statement of the theorem is proved. Next, consider the event

$$E_1 = \{\{x_n\}_{n \in \mathbb{Z}_{\geq 0}} \mid \exists \hat{x} \in S \setminus (F_1 \cap \dots \cap F_m) \text{ such that } \hat{x} \in \omega(x_n)\}.$$

Assume by contradiction that $\mathbb{P}[E_1] > 0$. Now we compute $\mathbb{P}[E|E_1]$. Note that, for each $\hat{x} \in S \setminus (F_1 \cap \dots \cap F_m)$, there exists a non-empty set $\mathcal{I}_{\hat{x}} \subset \{1, \dots, m\}$ such that, $\hat{x} = T_i(\hat{x})$ if $i \in \mathcal{I}_{\hat{x}}$, and $\hat{x} \neq T_i(\hat{x})$ if $i \notin \mathcal{I}_{\hat{x}}$. Similarly to the proof of Theorem 4.3, we can associate to each \hat{x} a positive real number δ such that the inequality $x \neq T_i(x)$ holds true for all $x \in B_\delta(\hat{x}) = \{x \in W \mid d(x, \hat{x}) \leq \delta\}$ and for all $i \notin \mathcal{I}_{\hat{x}}$. Moreover, we can define

$$\gamma_\delta = \min_{i \in \mathcal{I}_{\hat{x}}} \left\{ \min_{x \in B_\delta(\hat{x})} \left(U(x) - U(T_i(x)) \right) \right\},$$

where the continuity of the maps T_j , $j \in \{1, \dots, m\}$, and U , and the closedness of the set $B_\delta(\hat{x})$ ensure that $\gamma_\delta > 0$.

Consider the set $\{T_i\}_{i \in \mathcal{I}_{\hat{x}}}$; this is a collection of continuous maps having \hat{x} as fixed point. Therefore, there exists a suitable $\epsilon \in \mathbb{R}_{>0}$ such that, given any r -uple $(j_1, \dots, j_r) \in \mathcal{I}_{\hat{x}}^r$, $r \leq k$, we have $T_{j_1} \circ T_{j_2} \circ \dots \circ T_{j_r}(x) \in B_\delta(\hat{x})$ for all $x \in B_\epsilon(\hat{x})$. Given $\{x_n\}_{n \in \mathbb{Z}_{\geq 0}}$, if there exists $\hat{x} \in S \setminus (F_1 \cap \dots \cap F_m)$ such that $\hat{x} \in \omega(x_n)$, then there must exist $\{n_h \mid h \in \mathbb{Z}_{\geq 0}\}$ such that $x_{n_h} \in B_\epsilon(\hat{x})$ for all $h \in \mathbb{Z}_{\geq 0}$. Moreover, without loss of generality we can assume that $n_{h+1} - n_h > k$ for all $h \in \mathbb{Z}_{\geq 0}$. Consider now the event

$$E_3 = \{\{i_n \in \{1, \dots, m\} \mid n \in \mathbb{Z}_{\geq 0}\} \mid \exists \bar{h} \text{ such that } i_{n_h+s} \in \mathcal{I}_{\hat{x}} \text{ for all } s \in \{1, \dots, k\} \text{ and } h \geq \bar{h}\}.$$

To compute $\mathbb{P}[E_3]$, we define, for $j \in \mathbb{Z}_{\geq 0}$,

$$E_{3,j} = \{\{i_n \in \{1, \dots, m\} \mid n \in \mathbb{Z}_{\geq 0}\} \mid i_{n_h+s} \in \mathcal{I}_{\hat{x}} \text{ for all } s \in \{1, \dots, k\} \text{ and } h \geq j\}.$$

Observe that $\{E_{3,j} \mid j \in \mathbb{Z}_{\geq 0}\}$ is a countable collection of disjoint sets such that $E_3 = \bigcup_{j=0}^{\infty} E_{3,j}$. By hypothesis we have that

$$\mathbb{P}[E_{3,j}] \leq \lim_{l \rightarrow \infty} \prod_{s=j}^l (1-p) = 0,$$

and therefore $\mathbb{P}[E_3] = 0$. This implies that, almost surely, there exists a subsequence $\{n_{h_s}\}_{s \in \mathbb{Z}_{\geq 0}} \subseteq \{n_h\}_{h \in \mathbb{Z}_{\geq 0}}$ with the property that, for all $s \in \mathbb{Z}_{\geq 0}$, $x_{n_{h_s}+1} = T_i(x_{n_{h_s}})$ for some $i \notin \mathcal{I}_{\hat{x}}$ and, therefore, also with the property that $U(x_{n_{h_s}}) - U(x_{n_{h_s}+1}) > \gamma_\delta$.

Consequently, almost surely, we have that $\lim_{s \rightarrow \infty} U(x_{n_{h_s}}) = -\infty$ thus violating the fact that E is a certain event. This implies that $\mathbb{P}[E_1] = 0$ and that, almost surely, $x_n \rightarrow (F_1 \cap \dots \cap F_m) \cap U^{-1}(c)$. \square

REMARK 4.6. *The assumption, in Lemma 4.1 and Theorems 4.3 and 4.5, that W is strongly positively invariant ensures that any evolution of T with initial condition in W remains in W . By relaxing this assumption, it is possible to provide weaker versions of these results. Specifically, requiring W to be only compact and not necessarily strongly positively invariant, the thesis of Lemma 4.1 and Theorems 4.3 and 4.5 do not hold in general for any evolution of T with initial condition on W , but are still valid for those evolutions $\{x_n\}_{n \in \mathbb{Z}_{\geq 0}}$ that take values in W for all $n \in \mathbb{Z}_{\geq 0}$. \square*

5. On the topology of the space of partitions: compactness properties in the symmetric difference metric. Motivated by the invariance principles presented in Section 4, we study metric structures on the set of partitions, with a focus on compactness and continuity properties. Specifically, we show how a particular subset of the set of partitions can be regarded as a compact metric space and how certain relevant maps are continuous over that subspace. In this section, and only in this section, the assumptions on Q are relaxed to give more general results: we assume that $Q \subset \mathbb{R}^d$ is compact and connected and has non-empty interior.

Let \mathcal{C} denote the set of closed subsets of Q . We would like to introduce a topology on \mathcal{C} with two properties: \mathcal{C} is compact and the Voronoi map, the centroid map, and the multicenter function, defined in equations (2.2) (2.5), and (3.3) respectively, are continuous over \mathcal{C} . A natural candidate is the topology induced by the well-known [34] Hausdorff metric on \mathcal{C} : given two sets $A, B \in \mathcal{C}$, their Hausdorff distance is $d_H(A, B) = \max\{\max_{a \in A} \min_{b \in B} d(a, b), \max_{b \in B} \min_{a \in A} d(a, b)\}$. This metric induces a topology on \mathcal{C} which makes it a compact space, but is not suitable for our purpose because, with respect to this topology, the Voronoi map, the centroid map, and the multicenter function, are not continuous; see Appendix B. Additionally, note that, unlike the Hausdorff metric, the centroid map and the multicenter function are insensitive to sets of measure zero.

In what follows, we introduce the symmetric difference metric, as a metric insensitive to sets of measure zero. Given two subsets $A, B \in \mathcal{C}$, we define their *symmetric difference* by $A \ominus B = (A \cup B) \setminus (A \cap B)$. Moreover, letting μ denote the Lebesgue measure on \mathbb{R}^d , we define the *symmetric difference distance*, also called the *symmetric distance* for simplicity, $d_{\ominus} : \mathcal{C} \times \mathcal{C} \rightarrow \mathbb{R}_{\geq 0}$ by

$$d_{\ominus}(A, B) = \mu(A \ominus B),$$

that is, the symmetric distance between two sets is the measure of their symmetric difference. Given these notions, it is useful to identify sets that differ by a set of measure zero: we write $A \sim B$ whenever $\mu(A \ominus B) = 0$. Clearly, \sim is an equivalence relationship on \mathcal{C} and, accordingly, we let $\mathcal{C}^* = \mathcal{C}/\sim$ denote the *quotient set of closed subsets of Q* . Now, for any two elements A^* and B^* of \mathcal{C}^* , we define $d_{\ominus}(A^*, B^*) = d_{\ominus}(A, B)$ where A and B are any representatives of A^* and B^* , respectively. With this notion of d_{\ominus} on $\mathcal{C}^* \times \mathcal{C}^*$, it is easy to verify that $(\mathcal{C}^*, d_{\ominus})$ is a metric space. Unfortunately, it is known¹ that the metric space $(\mathcal{C}^*, d_{\ominus})$ is not compact. For this reason, we introduce a particular family of subsets of \mathcal{C} whose structure is sufficiently

¹For $Q = [0, 1]$, the sequence of sets $c_k = \cup_{i=1}^{2^k-1} [2(i-1)/2^k, (2i-1)/2^k]$, for $k \in \mathbb{N}$, satisfies $d_{\ominus}(c_i, c_j) = 1/2$ for all $i \neq j$. Hence, \mathcal{C} contains a sequence of $1/2$ -separated points with respect to the symmetric difference metric, violating compactness.

rich for our algorithm. For $\ell \in \mathbb{N}$, let $\mathcal{C}_{(\ell)} \subset \mathcal{C}$ denote the set of ℓ -convex and closed subsets of Q , that is, the set of subsets of Q equal to the union of ℓ convex and closed subsets of Q . Formally, we set

$$\mathcal{C}_{(\ell)} = \left\{ v \subseteq Q \mid v = \bigcup_{i=1}^{\ell} S_i \text{ where } S_1, \dots, S_{\ell} \subseteq Q \text{ are convex and closed} \right\}.$$

Note that we do not require the sets S_1, \dots, S_{ℓ} to be distinct so that $\mathcal{C}_{(k)} \subset \mathcal{C}_{(\ell)}$, for any $k < \ell$. In what follows we study the *quotient set of ℓ -convex and closed subsets* $\mathcal{C}_{(\ell)}^* = \mathcal{C}_{(\ell)} / \sim$. The next result is the main result of this section.

THEOREM 5.1 (Compactness of $\mathcal{C}_{(\ell)}^*$). *The pair $(\mathcal{C}_{(\ell)}^*, d_{\ominus})$ is a metric space and, with the topology induced by d_{\ominus} , the set $\mathcal{C}_{(\ell)}^*$ is compact.*

Proof. It is easy to verify that d_{\ominus} is a metric on $\mathcal{C}_{(\ell)}^*$. Instead, proving the compactness of $\mathcal{C}_{(\ell)}^*$ requires some attention. We aim to show that any sequence in $\mathcal{C}_{(\ell)}^*$ has a subsequence that converges to a point in $\mathcal{C}_{(\ell)}^*$. This fact's proof is articulated in several steps and relies upon several known results:

- (i) the space \mathcal{C} , endowed with the Hausdorff distance $d_H : \mathcal{C} \times \mathcal{C} \rightarrow \mathbb{R}_{\geq 0}$, is [31, Theorem 0.8] a compact metric space;
- (ii) if a sequence of closed convex subsets of Q converges in the Hausdorff sense to a set K , then [25, Proposition 1.6.8] K is closed and convex; and
- (iii) for any two convex subsets A, B of $Q \subset \mathbb{R}^d$, it is known [20, Eq. (1)] that

$$d_{\ominus}(A, B) \leq \left(\frac{2\kappa_d}{2^{1/d} - 1} \left(\frac{D}{2} \right)^{d-1} \right) d_H(A, B),$$

where $D = \max \{ \text{diam}(A), \text{diam}(B) \}$ and where κ_d is the volume of the unit sphere in \mathbb{R}^d .

Now let $\{v^*(n)\}_{n \in \mathbb{Z}_{\geq 0}}$ be any sequence in $\mathcal{C}_{(\ell)}^*$. For all $n \in \mathbb{Z}_{\geq 0}$, pick any representative of the equivalence class $v^*(n)$, denote it by $v(n) \in \mathcal{C}_{(\ell)}$ and consider the sequence $\{v(n)\}_{n \in \mathbb{Z}_{\geq 0}}$. Because $\{v(n)\}_{n \in \mathbb{Z}_{\geq 0}}$ is a sequence in $\mathcal{C}_{(\ell)}$ and because $\mathcal{C}_{(\ell)} \subseteq \mathcal{C}$, it follows from fact (i) above that $\{v(n)\}_{n \in \mathbb{Z}_{\geq 0}}$ contains a subsequence that converges in the Hausdorff sense to an element of \mathcal{C} . In other words, there exist $\{v(n_k)\}_{k \in \mathbb{Z}_{\geq 0}} \subseteq \{v(n)\}_{n \in \mathbb{Z}_{\geq 0}}$ and $\bar{v} \in \mathcal{C}$ such that $\lim_{k \rightarrow \infty} v(n_k) \stackrel{(H)}{=} \bar{v}$, where $\stackrel{(H)}{=}$ denotes convergence in the Hausdorff sense. We claim that $\bar{v} \in \mathcal{C}_{(\ell)}$ so that the set $\mathcal{C}_{(\ell)}$ is compact in the Hausdorff sense. To show this claim, we plan to exhibit a collection of convex and closed subsets of Q , say $\{S_1, \dots, S_{\ell}\}$, such that $\bar{v} = \bigcup_{i=1}^{\ell} S_i$. We begin the construction of \bar{v} as follows. By definition of $\mathcal{C}_{(\ell)}$, for all $k \in \mathbb{Z}_{\geq 0}$ there exists a collection $\{S_1(n_k), \dots, S_{\ell}(n_k)\}$ of convex and closed subsets of Q whose union equals $v(n_k)$. Now, we consider the sequence $\{S_1(n_k)\}_{k \in \mathbb{Z}_{\geq 0}}$. Again, since (\mathcal{C}, d_H) is a compact metric space we have that there exists a subsequence $\{n_{k_1}\}_{k_1 \in \mathbb{Z}_{\geq 0}} \subseteq \{n_k\}_{k \in \mathbb{Z}_{\geq 0}}$ such that $\lim_{k_1 \rightarrow \infty} S_1(n_{k_1}) \stackrel{(H)}{=} \bar{S}_1$ for some \bar{S}_1 . Fact (ii) above ensures that \bar{S}_1 is a convex closed subset of Q . Consider now the sequence $\{S_2(n_{k_1})\}_{k_1 \in \mathbb{Z}_{\geq 0}}$. By reasoning as previously we know that there exists a subsequence $\{n_{k_2}\}_{k_2 \in \mathbb{Z}_{\geq 0}} \subseteq \{n_{k_1}\}_{k_1 \in \mathbb{Z}_{\geq 0}}$ such that $\lim_{k_2 \rightarrow \infty} S_2(n_{k_2}) \stackrel{(H)}{=} \bar{S}_2$ where \bar{S}_2 is some convex closed set of Q . Moreover, it is clear that also $\lim_{k_2 \rightarrow \infty} S_1(n_{k_2}) \stackrel{(H)}{=} \bar{S}_1$. By iterating this procedure we conclude that there exist a sequence $\{n_s\}_{s \in \mathbb{Z}_{\geq 0}}$ and a collection of convex and closed sets $\{\bar{S}_1, \dots, \bar{S}_{\ell}\}$ such that $\lim_{s \rightarrow \infty} S_i(n_s) \stackrel{(H)}{=} \bar{S}_i$ for all $i \in \{1, \dots, \ell\}$. Next, we aim to show that $v(n_s)$ converges to $\bigcup_{i=1}^{\ell} \bar{S}_i$ in the Hausdorff sense, that is,

$$(5.1) \quad \lim_{s \rightarrow \infty} \bigcup_{i=1}^{\ell} S_i(n_s) = \bigcup_{i=1}^{\ell} \lim_{s \rightarrow \infty} S_i(n_s) = \bigcup_{i=1}^{\ell} \bar{S}_i.$$

For simplicity, let us denote $\lim_{s \rightarrow \infty} \bigcup_{i=1}^{\ell} S_i(n_s)$ by S_{∞} . For $p \in Q$, note

$$p \in S_{\infty} \iff \text{dist} \left(p, \lim_{s \rightarrow \infty} \bigcup_{i=1}^{\ell} S_i(n_s) \right) = 0 \iff \lim_{s \rightarrow \infty} \text{dist} \left(p, \bigcup_{i=1}^{\ell} S_i(n_s) \right) = 0,$$

where, for a given closed set X , $\text{dist}(p, X)$ denotes the Euclidean distance between p and X , namely, $\min_{x \in X} \|p - x\|$. Using the fact that, for given closed sets X and Y , $\text{dist}(p, X \cup Y) = \min \{ \text{dist}(p, X), \text{dist}(p, Y) \}$, we can write

$$\begin{aligned} \lim_{s \rightarrow \infty} \text{dist} \left(p, \bigcup_{i=1}^{\ell} S_i(n_s) \right) = 0 &\iff \lim_{s \rightarrow \infty} \min \{ \text{dist}(p, S_1(n_s)), \dots, \text{dist}(p, S_{\ell}(n_s)) \} = 0 \\ &\iff \exists j \in \{1, \dots, \ell\} \text{ s.t. } \lim_{s \rightarrow \infty} \text{dist}(p, S_j(n_s)) = 0 \\ &\iff \exists j \in \{1, \dots, \ell\} \text{ s.t. } p \in \bar{S}_j \\ &\iff p \in \bigcup_{i=1}^{\ell} \bar{S}_i. \end{aligned}$$

The above chain of implications proves (5.1). Now observe that, from the uniqueness of the limit it follows that $\lim_{k \rightarrow \infty} v(n_k) = \lim_{s \rightarrow \infty} \bigcup_{i=1}^{\ell} S_i(n_s) = \bigcup_{i=1}^{\ell} \bar{S}_i = \bar{v}$. Since \bar{S}_i is closed and convex for all $i \in \{1, \dots, \ell\}$, it follows that $\bar{v} \in \mathcal{C}_{(\ell)}$ and, in turn, that $\mathcal{C}_{(\ell)}$ endowed with the Hausdorff metric is a compact metric space.

To establish the statement of the Theorem we still need to prove that $\lim_{k \rightarrow \infty} d_{\ominus}(v(n_k), \bar{v}) = 0$, or that $\lim_{s \rightarrow \infty} d_{\ominus}(v(n_s), \bar{v}) = 0$. To this end, observe that, given $X_1, X_2, Y_1, Y_2 \subseteq Q$, the following inclusion holds $(X_1 \cup X_2) \ominus (Y_1 \cup Y_2) \subseteq (X_1 \ominus Y_1) \cup (X_2 \ominus Y_2)$. Hence, we compute

$$v(n_s) \ominus \bar{v} = \left(\bigcup_{i=1}^{\ell} S_i(n_s) \right) \ominus \bar{v} = \left(\bigcup_{i=1}^{\ell} S_i(n_s) \right) \ominus \left(\bigcup_{i=1}^{\ell} \bar{S}_i \right) \subseteq \bigcup_{i=1}^{\ell} (S_i(n_s) \ominus \bar{S}_i),$$

which implies

$$\begin{aligned} d_{\ominus}(v(n_s), \bar{v}) &= d_{\ominus} \left(\bigcup_{i=1}^{\ell} S_i(n_s), \bar{v} \right) \leq \sum_{i=1}^{\ell} d_{\ominus}(S_i(n_s), \bar{S}_i) \\ &\leq \sum_{i=1}^{\ell} \frac{2\kappa_d}{2^{1/d} - 1} \left(\frac{\max \{ \text{diam}(S_i(n_s)), \text{diam}(\bar{S}_i) \}}{2} \right)^{d-1} d_H(S_i(n_s), \bar{S}_i) \\ &\leq \frac{2\kappa_d}{2^{1/d} - 1} \left(\frac{\text{diam}(Q)}{2} \right)^{d-1} \sum_{i=1}^{\ell} d_H(S_i(n_s), \bar{S}_i), \end{aligned}$$

where the second and the third inequalities follow, respectively, from fact (iii) above and from the upper bounds $\max \{ \text{diam}(S_i(n_s)), \text{diam}(\bar{S}_i) \} \leq \text{diam}(Q)$, for all $i \in \{1, \dots, \ell\}$. Since $\lim_{s \rightarrow \infty} d_H(S_i(n_s), \bar{S}_i) = 0$ for all $i \in \{1, \dots, \ell\}$, we conclude that

$$\lim_{s \rightarrow \infty} d_{\ominus}(v(n_s), \bar{v}) = 0.$$

Now, let \bar{v}^* denote the equivalence class for which \bar{v} is a representative. Since the metric d_{\ominus} is insensitive to sets of measure zero, it follows that $\lim_{s \rightarrow \infty} d_{\ominus}(v^*(n_s), \bar{v}^*) = 0$

and, in turn, that $\{v^*(n)\}_{n \in \mathbb{Z}_{\geq 0}}$ has a subsequence convergent to point of $\mathcal{C}_{(\ell)}^*$. This concludes the proof that $\mathcal{C}_{(\ell)}^*$ is a compact space. \square

The metric d_{\ominus} naturally extends to a metric over the space $(\mathcal{C}^*)^N$, and hence over $(\mathcal{C}_{(\ell)}^*)^N$, by defining

$$(5.2) \quad d_{\ominus}(u, v) = \sum_{i=1}^N d_{\ominus}(u_i, v_i),$$

for any $u = (u_i)_{i=1}^N$ and $v = (v_i)_{i=1}^N$ in $(\mathcal{C}^*)^N$. The compactness of $(\mathcal{C}_{(\ell)}^*)^N$ is then a simple consequence of Theorem 5.1.

COROLLARY 5.2 (Compactness of $(\mathcal{C}_{(\ell)}^*)^N$). *The pair $((\mathcal{C}_{(\ell)}^*)^N, d_{\ominus})$ is a metric space and, with the topology induced by d_{\ominus} , $(\mathcal{C}_{(\ell)}^*)^N$ is compact.*

6. On the continuity of some geometric maps and the resulting convergence proofs. In this section we prove the main convergence theorems for our gossip coverage algorithms. First, however, we need to establish the continuity properties of certain geometric maps and of the proposed algorithms T and T^{δ} . Before proceeding, we discuss two significant modeling aspects. First, recall that Section 5 introduces the spaces \mathcal{C} , $\mathcal{C}_{(\ell)}$, \mathcal{C}^N , $\mathcal{C}_{(\ell)}^N$ and the corresponding quotient sets \mathcal{C}^* , $\mathcal{C}_{(\ell)}^*$, $(\mathcal{C}^*)^N$, $(\mathcal{C}_{(\ell)}^*)^N$. We can do the same with the partition space \mathcal{V}_N . Indeed from Definition 2.1(iii) we have that each component v_i of $v \in \mathcal{V}_N$ can be mapped by the canonical projection into an equivalence class v_i^* in $\mathcal{C}^* \setminus \{\emptyset\}$; in turn, any $v \in \mathcal{V}_N$ can be naturally associated to the N -collection of equivalence classes $v^* = (v_i^*)_{i=1}^N$. Accordingly, we denote the *space of equivalence classes of N -partitions* by \mathcal{V}_N^* . Recall that all these quotient spaces are metric spaces with respect to the symmetric distance d_{\ominus} .

Second, recall the following maps: the centroid map $\text{Cd} : \{A \in \mathcal{C} \mid \mu(A) > 0\} \rightarrow Q$ defined in equation (2.5), the 1-center function $\mathcal{H}_1 : Q \times \mathcal{C} \rightarrow \mathbb{R}_{\geq 0}$ defined in equation (2.4), the multicenter function $\mathcal{H}_{\text{centroid}} : \mathcal{V}_N \rightarrow \mathbb{R}_{\geq 0}$ defined in equation (3.3), the gossip coverage map $T_{ij} : \mathcal{V}_N \rightarrow \mathcal{V}_N$ with $i \neq j$ defined in equation (3.2), and, for $\delta > 0$ and $i \neq j$, the modified gossip coverage map $T_{ij}^{\delta} : \mathcal{V}_N \rightarrow \mathcal{V}_N$ defined in Section 3.4. We claim that all these maps are insensitive to sets of measure zero. To substantiate this claim, observe that the integrals of a bounded measurable function over a set A and over a set B are equal if $d_{\ominus}(A, B) = 0$. This observation allows us to redefine the centroid map, the 1-center function and the multicenter function as $\text{Cd} : \mathcal{C}^* \setminus \{\emptyset\} \rightarrow Q$, $\mathcal{H}_1 : Q \times \mathcal{C}^* \rightarrow \mathbb{R}_{\geq 0}$ and $\mathcal{H}_{\text{centroid}} : \mathcal{V}_N^* \rightarrow \mathbb{R}_{\geq 0}$, respectively. Regarding the modified gossip coverage map T_{ij}^{δ} , we reason as follows. For $v^* \in \mathcal{V}_N^*$, let $v \in \mathcal{V}_N$ and $v' \in \mathcal{V}_N$ be two representatives of v^* and let \hat{v} and \hat{v}' denote, respectively, the images of v and v' under the map T_{ij}^{δ} , that is, $\hat{v} = T_{ij}^{\delta}(v)$ and $\hat{v}' = T_{ij}^{\delta}(v')$. Since the centroid map and the definitions of the points \hat{p}_i and \hat{p}_j in equation (3.5) are insensitive to sets of measure zero, it follows that $d_{\ominus}(\hat{v}, \hat{v}') = 0$; in other words \hat{v} and \hat{v}' belong to the same equivalence class, say $\hat{v}^* \in \mathcal{V}_N^*$. From these facts we can redefine the modified gossip coverage map as $T_{ij}^{\delta} : \mathcal{V}_N^* \rightarrow \mathcal{V}_N^*$. An analogous argument applies to the map T . This concludes the justification of our claim. Finally, note that the Voronoi map $V : Q^N \setminus S_N \rightarrow \mathcal{V}_N$ defined in equation (2.2) can be composed with the standard quotient projection map and therefore denoted by $V : Q^N \setminus S_N \rightarrow \mathcal{V}_N^*$. In summary, the centroid map, the 1-center function, the multicenter function, the modified gossip coverage map, and the Voronoi map are indeed insensitive to sets of measure zero.

6.1. Continuity of various geometric maps. We start by recalling that the compact connected set Q is equipped with a bounded measurable positive function $\phi : Q \rightarrow \mathbb{R}_{>0}$. We define the diameter of Q and the infinity norm of ϕ by $\text{diam}(Q) = \max\{\|x - y\| \mid x, y \in Q\}$ and $\|\phi\|_\infty = \max_{x \in Q} \phi(x)$, respectively. The following lemma states some important properties of the 1-center cost function.

LEMMA 6.1 (Continuity properties of the 1-center function). *Given a compact convex set $Q \subset \mathbb{R}^d$, let $\phi : Q \rightarrow \mathbb{R}_{>0}$ be bounded and measurable and let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ be locally Lipschitz, increasing, and convex. Define the function $\mathcal{H}_1 : Q \times \mathcal{C}^* \rightarrow \mathbb{R}_{\geq 0}$ as in equation (2.4). The following statements hold:*

- (i) *the function $p \mapsto \mathcal{H}_1(p; A)$ is strictly convex in p , for any $A \in \mathcal{C}^* \setminus \{\emptyset\}$,*
- (ii) *the function $p \mapsto \mathcal{H}_1(p; A)$ is globally Lipschitz in p , for any $A \in \mathcal{C}^*$, and*
- (iii) *the function $A \mapsto \mathcal{H}_1(p; A)$ is globally Lipschitz in A with respect to d_\ominus , for any $p \in Q$.*

We now state the main result of this subsection.

THEOREM 6.2 (Continuity properties of centroid, multicenter and Voronoi maps). *Given a compact convex set $Q \subset \mathbb{R}^d$, let $\phi : Q \rightarrow \mathbb{R}_{>0}$ be bounded and measurable and let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ be locally Lipschitz, increasing, and convex. With respect to the topology induced by d_\ominus , the following maps are continuous:*

- (i) *the centroid map $\text{Cd} : \mathcal{C}^* \setminus \{\emptyset\} \rightarrow Q$,*
- (ii) *the multicenter function $\mathcal{H}_{\text{centroid}} : \mathcal{V}_N^* \rightarrow \mathbb{R}_{\geq 0}$,*
- (iii) *the Voronoi map $V : Q^N \setminus S_N \rightarrow \mathcal{V}_N^*$,*
- (iv) *for all $i, j \in \{1, \dots, N\}$, $i \neq j$, the gossip coverage map $T_{ij} : \{v \in \mathcal{V}_N^* \mid \text{Cd}(v_i) \neq \text{Cd}(v_j)\} \rightarrow \mathcal{V}_N^*$, and*
- (v) *for all $\delta > 0$, $i, j \in \{1, \dots, N\}$, $i \neq j$, the modified gossip coverage map $T_{ij}^\delta : \mathcal{V}_N^* \rightarrow \mathcal{V}_N^*$.*

The continuity properties (ii) and (iv) (respectively, (v)) are exactly what is needed to apply the Krasovskii-LaSalle invariance principles stated in Section 4 to the gossip coverage algorithm (respectively, to the modified gossip coverage algorithm). The continuity properties (i) and (iii) are intermediate results of independent interest.

Proof. [Proof of Lemma 6.1] Let L_f be the Lipschitz constant of $f : [0, \text{diam}(Q)] \rightarrow \mathbb{R}_{\geq 0}$. We check the claims in order, beginning with statement (i). For $\lambda \in (0, 1)$, $A \in \mathcal{C}^* \setminus \{\emptyset\}$, and $p_1, p_2 \in Q$, we compute

$$\begin{aligned}
 \mathcal{H}_1(\lambda p_1 + (1 - \lambda)p_2; A) &= \int_A f(\|\lambda p_1 + (1 - \lambda)p_2 - q\|)\phi(q) dq \\
 (6.1) \quad &\leq \int_A f(\lambda\|p_1 - q\| + (1 - \lambda)\|p_2 - q\|)\phi(q) dq \\
 (6.2) \quad &\leq \int_A \left(\lambda f(\|p_1 - q\|) + (1 - \lambda)f(\|p_2 - q\|) \right) \phi(q) dq \\
 &= \lambda \mathcal{H}_1(p_1; A) + (1 - \lambda) \mathcal{H}_1(p_2; A),
 \end{aligned}$$

where inequality (6.1) follows from the triangle inequality and from f being increasing, and inequality (6.2) follows from the convexity of f . This inequality proves convexity. Moreover, since the first inequality is strict outside the line passing through p_1 and p_2 and since A has non-empty interior, the function is in fact strictly convex. Note that statement (i) implies that $p \mapsto \mathcal{H}_1(p; A)$ is locally Lipschitz, using [33, Theorem 10.4].

The stronger statement (ii) can be derived as follows. For $p_1, p_2 \in Q$, we compute

$$\begin{aligned}
|\mathcal{H}_1(p_1; A) - \mathcal{H}_1(p_2; A)| &= \left| \int_A f(\|p_1 - q\|)\phi(q) dq - \int_A f(\|p_2 - q\|)\phi(q) dq \right| \\
&= \left| \int_A [f(\|p_1 - q\|) - f(\|p_2 - q\|)]\phi(q) dq \right| \\
&\leq \int_A |f(\|p_1 - q\|) - f(\|p_2 - q\|)|\phi(q) dq \\
&\leq \int_A L_f \|p_1 - p_2\| \phi(q) dq \leq L_f \|\phi\|_\infty \mu(A) \|p_1 - p_2\|.
\end{aligned}$$

This implies the Lipschitz condition in statement (ii). Statement (iii) can be proved as follows. Let A, A' be two elements of \mathcal{C}^* , note $A \ominus A' = (A \setminus A') \cup (A \cap A')$ and compute

$$\begin{aligned}
|\mathcal{H}_1(p; A) - \mathcal{H}_1(p; A')| &= \left| \int_{A \setminus A'} f(\|p - q\|)\phi(q) dq - \int_{A' \setminus A} f(\|p - q\|)\phi(q) dq \right| \\
&\leq \left| \int_{A \setminus A'} f(\|p - q\|)\phi(q) dq \right| + \left| \int_{A' \setminus A} f(\|p - q\|)\phi(q) dq \right| \\
&= \int_{A \ominus A'} f(\|p - q\|)\phi(q) dq \\
&\leq \max\{f(\|p - q\|) \mid p, q \in A \ominus A'\} \|\phi\|_\infty \mu(A \ominus A') \\
&\leq f(\text{diam}(Q)) \|\phi\|_\infty d_\ominus(A, A'),
\end{aligned}$$

where last inequality follows from f being increasing. The bound implies the Lipschitz condition. \square

Before proving Theorem 6.2 we need the following lemma about perturbations of convex optimization problems.

LEMMA 6.3. *Given a compact convex set $Q \subset \mathbb{R}^d$ and a metric space (X, d) , let $H : Q \times X \rightarrow \mathbb{R}$ have the properties that*

- (i) *the map $x \mapsto H(q, x)$ is globally Lipschitz for all $q \in Q$, and*
- (ii) *the map $q \mapsto H(q, x)$ is continuous and strictly convex.*

Then the map $q^ : X \rightarrow Q$, defined by $q^*(x) = \text{argmin}_{q \in Q} H(q, x)$, is continuous.*

Proof. Let L_H be the Lipschitz constant of $x \mapsto H(q, x)$. Thanks to the Lipschitz condition of the function $x \mapsto H(q, x)$, for all $x, y \in X$, the point of minimum $q^*(x)$ takes value in $S = \{q \in Q \mid H(q, y) \leq H(q^*(y), y) + 2L_H d(y, x)\}$. Since S is a sub-level set of the strictly convex function $q \mapsto H(q, y)$, and since the diameter of a sub-level set depends continuously on the level, the distance $\|q^*(y) - q^*(x)\|$ can be made arbitrary small by reducing $d(y, x)$. This implies the claimed continuity. \square

We are now ready to prove the main result.

Proof. [Proof of Theorem 6.2] We prove the theorem claims in the order in which they are presented. Claim (i) follows combining Lemma 6.3 and Lemma 6.1. Since the multicenter function $\mathcal{H}_{\text{centroid}}$ is a sum of suitable 1-center functions \mathcal{H}_1 , the claim (ii) is also immediate.

Regarding claim (iii), we begin by discussing in detail the two dimensional case. Let $N = 2$, and p_1 and p_2 be two points in Q . Let $2l = \|p_1 - p_2\|$. Since $p_1 \neq p_2$, $l > 0$. Up to isometries, we can assume that, in the Euclidean plane (x, y) , $p_1 = (-l, 0)$ and $p_2 = (l, 0)$. Let d_1 and d_2 be the distances from the origin of points p_1 and p_2 , respectively. It is clear that the two Voronoi regions of p_1 and p_2 are separated by

the locus of points $\{x \in Q \mid \|x - p_1\| = \|x - p_2\|\}$, that is the vertical axis. Now, we assume that the positions of p_1 and p_2 are perturbed by a quantity less than or equal to ϵ , with $0 < \epsilon < l$. By effect of the perturbation, the axis separating the two Voronoi regions is perturbed, but it is contained in the locus of points $Y_{12}(\epsilon) = \{x \in Q \mid \|x - p_1\| - \|x - p_2\| \leq 2\epsilon\}$. By definition, this is the set comprised between the two branches of the hyperbola whose equation is $\frac{x^2}{\epsilon^2} - \frac{y^2}{l^2 - \epsilon^2} = 1$. By elementary geometric considerations, the area of this region can be upper bounded by

$$\mu(Y_{12}(\epsilon)) \leq 2\epsilon \operatorname{diam}(Q) + 4 \operatorname{diam}(Q)^2 \frac{\epsilon/l}{\sqrt{1 - \frac{\epsilon^2}{l^2}}} \leq 4 \operatorname{diam}(Q) \left(1 + \frac{\operatorname{diam}(Q)}{l}\right) \epsilon.$$

This bound implies the continuity. The case in which $N > 2$ follows because, moving all points by at most ϵ , the change in all the regions is upper bounded by $\bigcup_{1 \leq i, j \leq N} Y_{ij}(\epsilon)$, which vanishes as $\epsilon \rightarrow 0^+$.

The last remaining step is to prove claims (iv) and (v). We focus on claim (v) and show claim (iv) as a byproduct. Let $v \in \mathcal{V}_N$ and let $\hat{v} = T_{ij}^\delta(v)$. According to the definitions in Subsection 3.4, \hat{v} is characterized by the sets \hat{R}_i and \hat{R}_j . Recall that these two sets depend on the sets R_i, R_j , on the scalar $\beta_{ij}(v)$ and on the points \hat{p}_i and \hat{p}_j . One can see that $\beta_{ij}(v)$ is a continuous function of its arguments v_i and v_j . Hence, it suffices to show that also \hat{R}_i, \hat{R}_j and \hat{p}_i, \hat{p}_j depend continuously on v_i and v_j . To do this, introduce $v' \in \mathcal{V}_N$ and compute $\hat{v}' = T_{ij}^\delta(v')$. Assume $\operatorname{Cd}(v_i) \neq \operatorname{Cd}(v_j)$ and $\operatorname{Cd}(v'_i) \neq \operatorname{Cd}(v'_j)$. Analogously to how we defined R_i and R_j , we now define the regions $R'_i = v'_i \cap H_{\text{bisector}}(\operatorname{Cd}(v'_j), \operatorname{Cd}(v'_i))$ and $R'_j = v'_j \cap H_{\text{bisector}}(\operatorname{Cd}(v'_i), \operatorname{Cd}(v'_j))$. We aim to upper bound the composite distance $d_\ominus(R_i, R'_i) + d_\ominus(R_j, R'_j)$. Observe that this composite distance depends on the difference of the two argument regions v_i, v'_i and v_j, v'_j both directly and indirectly via the induced difference between the centroids. Recalling the proof of claim (iii), let ϵ be an upper bound on the displacement between the two centroids. Then the region $Y(\epsilon) = \{x \in Q \mid \|\|x - \operatorname{Cd}(v_i)\| - \|x - \operatorname{Cd}(v_j)\|\| \leq 2\epsilon\}$ needs to be included in the upper bound on the composite distance. Combining these considerations we obtain

$$(6.3) \quad d_\ominus(R_i, R'_i) + d_\ominus(R_j, R'_j) \leq (d_\ominus(v_i, v'_i) + d_\ominus(v_j, v'_j)) \\ + \mu(Y(\max\{\|\operatorname{Cd}(v_i) - \operatorname{Cd}(v'_i)\|, \|\operatorname{Cd}(v_j) - \operatorname{Cd}(v'_j)\|\})).$$

Clearly, if $d_\ominus(v_i, v'_i) \rightarrow 0$ and $d_\ominus(v_j, v'_j) \rightarrow 0$, then $\|\operatorname{Cd}(v_i) - \operatorname{Cd}(v'_i)\| \rightarrow 0$ and $\|\operatorname{Cd}(v_j) - \operatorname{Cd}(v'_j)\| \rightarrow 0$ and, in turn, also $d_\ominus(R_i, R'_i) + d_\ominus(R_j, R'_j) \rightarrow 0$. Hence, we can argue that the sets R_i and R_j depend continuously on the regions v_i and v_j . This is enough to prove statement (iv), provided the two regions v_i and v_j have distinct centroids. Moreover a direct consequence of this fact is that also the points \hat{p}_i and \hat{p}_j depend continuously on v_i and v_j . Finally, observe that in the limit case $\operatorname{Cd}(v_i) = \operatorname{Cd}(v_j)$ the continuity of R_i and R_j is captured by the fact that $\beta_{ij}(v)$ is a continuous function of $\operatorname{Cd}(v_i), \operatorname{Cd}(v_j)$ and that $\beta_{ij}(v) = 0$ if $\operatorname{Cd}(v_i) = \operatorname{Cd}(v_j)$. The continuity of $R_i, R_j, \beta_{ij}(v), \hat{p}_i$ and \hat{p}_j imply also the continuity of \hat{R}_i and \hat{R}_j and, in turn, of T_{ij}^δ . \square

6.2. Convergence proofs. In view of the identification between N -partitions and their equivalence classes introduced at the beginning of this section, we are now ready to complete the proof of the convergence results presented in Section 3.3.

We start by clarifying the precise meaning of convergence in Theorems 3.7 and 3.8. Specifically, we say that a sequence of partitions $\{v(t)\}_{t \in \mathbb{Z}_{\geq 0}} \subset \mathcal{V}_N$ converges to a set

of partitions $X \subset \mathcal{V}_N$ if the symmetric distance from $\{v(t)\}_{t \in \mathbb{Z}_{\geq 0}}$ to X converges to zero, that is,

$$\lim_{t \rightarrow \infty} \inf \{d_{\ominus}(v(t), x) \mid x \in X\} = 0.$$

Proof. [Proof of Theorem 3.8] We prove the deterministic statement (i). We start by observing that, through the canonical projection, the evolution $\{v(t)\}_{t \in \mathbb{Z}_{\geq 0}} \subset \mathcal{V}_N$ of $T^{\delta} : \mathcal{V}_N \rightarrow \mathcal{V}_N$ can be mapped into the evolution $\{v^*(t)\}_{t \in \mathbb{Z}_{\geq 0}} \subset \mathcal{V}_N^*$ of $T^{\delta} : \mathcal{V}_N^* \rightarrow \mathcal{V}_N^*$. We aim to apply Theorem 4.3 to the dynamical system $T^{\delta} : \mathcal{V}_N^* \rightarrow \mathcal{V}_N^*$ and its evolution $\{v^*(t)\}_{t \in \mathbb{Z}_{\geq 0}} \subset \mathcal{V}_N^*$. In what follows, our goal is to verify whether Assumptions (i), (ii), (iii) and (iv) of Theorem 4.3 are satisfied.

Since $\{v(t)\}_{t \in \mathbb{Z}_{\geq 0}}$ is non-vanishing and finitely convex by assumption, it follows that there exists $\ell \in \mathbb{N}$ such that the ω -limit set of $\{v(t)\}_{t \in \mathbb{Z}_{\geq 0}}$ is contained in $\mathcal{C}_{(\ell)}^N \cap \mathcal{V}_N$, that is, $\omega(v(t)) \subseteq \mathcal{C}_{(\ell)}^N \cap \mathcal{V}_N$. This implies also that $\omega(v^*(t)) \subseteq (\mathcal{C}_{(\ell)}^*)^N \cap \mathcal{V}_N^*$. As stated in Theorem 5.1, $\mathcal{C}_{(\ell)}^*$ is compact in the topology induced by the metric d_{\ominus} . Hence, even though $(\mathcal{C}_{(\ell)}^*)^N$ is not strongly positive invariant for T^{δ} , the weaker version of Assumption (i) of Theorem 4.3, as given in Remark 4.6, holds true for the sequence $\{v^*(t)\}_{t \in \mathbb{Z}_{\geq 0}} \subset \mathcal{V}_N^*$. Now, as one can deduce from Theorem 6.2(ii) and Lemma 3.4, the function $\mathcal{H}_{\text{centroid}}$ satisfies the Assumption (ii) of Theorem 4.3, thus playing the role of a Lyapunov function for the dynamical system T^{δ} . Moreover, from Theorem 6.2(v), note that the system evolves through maps that are continuous in \mathcal{V}_N^* with respect to the metric d_{\ominus} : thus the Assumption (iii) of Theorem 4.3 is satisfied. Finally observe that the Assumption (iv) of Theorem 4.3 corresponds to the assumption of uniform persistency in Theorem 3.8. Therefore, we conclude that the evolution $\{v^*(t)\}_{t \in \mathbb{Z}_{\geq 0}}$ converges to the intersection of the fixed points of the maps T_{ij}^{δ} , for all $i, j \in \{1, \dots, N\}$, $j \neq i$. According to Lemma 3.3, this intersection coincides with the set of mixed centroidal Voronoi partitions up to sets of measure zero.

The proof of the stochastic statement (ii) follows the same lines, applying Theorem 4.5 instead of Theorem 4.3. \square

Proof. [Proof of Theorem 3.7] The proof of this result follows the lines of the proof above with two distinctions. The distinctions come from the assumption that the evolution v , in addition to being non-vanishing and finitely convex, is also distinct centroidal. The first distinction is as follows. In order to apply the Krasovskii-LaSalle invariance principle we require the continuity property stated in Theorem 6.2(iv). Additionally, we note that the space of finitely convex and distinct centroidal partitions

$$\{v \in \mathcal{V}_N \cap \mathcal{C}_{(\ell)}^N \mid \|\text{Cd}(v_i) - \text{Cd}(v_j)\| \geq \epsilon \text{ for all } i \neq j\}$$

is a closed and hence compact subspace of \mathcal{V}_N . The second distinction is as follows. Since we rule out the case of coincident centroids, we can infer convergence to centroidal Voronoi partitions instead of convergence to mixed centroidal Voronoi partitions; see Lemma 3.3(ii). \square

7. Conclusions. In summary, we have introduced novel coverage, deployment and partitioning algorithms for robotic networks with minimal communication requirements. To analyze our proposed algorithms, we have developed and characterized (1) intuitive versions of the Krasovskii-LaSalle Invariance Principle for deterministic and stochastic switching systems, (2) relevant topological properties of the space of partitions, and (3) useful continuity properties of a number of geometric and multi-center functions.

We believe there remain interesting open issues in the study of gossiping robots and of dynamical systems on the space of partitions. We are keen on extending these ideas to non-convex complex environments and discrete environments such as graphs; see [17] for some preliminary work in this direction. Following Remark 3.2 we plan to study gossip coverage algorithms for more general multicenter functions, including nonsmooth, anisotropic and inhomogeneous functions. Additionally, we plan to investigate gossip coverage algorithms capable of adapting to time-varying scenarios such as problems in which robotic agents arrive to and depart from the network. Finally, inspired by stigmergy in territorial animals, we plan to design communication protocols for multiagent systems based on the ability of leaving messages in the environment.

Appendix A. A counterexample showing the necessity of uniformly persistent switches. Theorem 4.3 contains a persistent switching conditions, that is, it requires the existence of $D \in \mathbb{N}$ such that every map T_i , $i \in \{1, \dots, m\}$, is applied at least once within every interval $[n, n + D]$, $n \in \mathbb{Z}_{\geq 0}$. This appendix contains an example proving the necessity of this condition.

Consider the plane in polar coordinates $X = \mathbb{R}_{>0} \times [0, 2\pi[\cup \{0, 0\}$. Define the standard metric $d : X \times X \rightarrow \mathbb{R}_{\geq 0}$ as follows: let (ρ_1, θ_1) , (ρ_2, θ_2) be any pair of elements of X and

$$d((\rho_1, \theta_1), (\rho_2, \theta_2)) = \sqrt{(\rho_1 \cos \theta_1 - \rho_2 \cos \theta_2)^2 + (\rho_1 \sin \theta_1 - \rho_2 \sin \theta_2)^2}.$$

Consider now the continuous maps $T_i : X \rightarrow X$, $i \in \{1, 2\}$, defined by respectively

$$T_1(\rho, \theta) = \begin{cases} (\rho^2, \theta), & \text{if } 0 \leq \rho \leq 1, \\ \left(\frac{2\rho-1}{\rho}, (\theta + \rho - 1) \bmod 2\pi \right), & \text{if } \rho > 1, \end{cases}$$

$$T_2(\rho, \theta) = \begin{cases} ((1 - \sin \theta)\rho, \theta), & \text{if } 0 \leq \theta \leq \pi, \\ (\rho, \theta) & \text{if } \pi \leq \theta \leq 2\pi. \end{cases}$$

Define $T : X \rightrightarrows X$ by $T(\rho, \theta) = \{T_1(\rho, \theta), T_2(\rho, \theta)\}$ and the function $U : X \rightarrow \mathbb{R}_{\geq 0}$ by $U(\rho, \theta) = \rho$. Observe that U is continuous and non-increasing along T . Assume now that there exists $D \in \mathbb{N}$ such that, for any $n \in \mathbb{Z}_{\geq 0}$, there exist n_1 and n_2 within the interval $]n, n + D]$ such that $x_{n_1+1} = T_1(x_{n_1})$ and $x_{n_2+1} = T_2(x_{n_2})$. Then, by Theorem 4.3, the ω -limit set of each evolution of T is a subset of

$$(A.1) \quad \{(\rho, \theta) \in X \mid \rho = 1, \pi \leq \theta \leq 2\pi\} \cup \{0, 0\}.$$

Next, we relax the condition that the map T_2 is applied at least once inside each interval of arbitrary amplitude D and we show that there exists one sequence that does not converge to the ω -limit set in equation (A.1). To this aim, assume the sequence $\{(\rho(n), \theta(n))\}_{n \in \mathbb{Z}_{\geq 0}}$ satisfies

- (i) $\rho(0) > 1$;
- (ii) $(\rho(1), \theta(1)) = T_1(\rho(0), \theta(0))$ and
- (iii) $(\rho(n+1), \theta(n+1)) = T_2(\rho(n), \theta(n))$ if and only if $\pi \leq \theta(n) \leq 2\pi$ and $(\rho(n), \theta(n)) = T_1(\rho(n-1), \theta(n-1))$.

Note that if $\pi \leq \theta(n) \leq 2\pi$, then $T_2(\rho(n), \theta(n)) = (\rho(n), \theta(n))$. Therefore, the evolution $\{(\rho(n), \theta(n))\}$ equals $\{(\hat{\rho}(n), \hat{\theta}(n))\}$ where $(\hat{\rho}(0), \hat{\theta}(0)) = (\rho(0), \theta(0))$ and

$(\widehat{\rho}(n), \widehat{\theta}(n)) = T_1^n (\widehat{\rho}(0), \widehat{\theta}(0))$. Regarding this new sequence, observe that

$$(A.2) \quad 1 < \widehat{\rho}(i) < 2 \text{ and } \widehat{\rho}(i+1) < \widehat{\rho}(i), \text{ for all } i \geq 1,$$

$$(A.3) \quad 0 < \widehat{\theta}(i+1) - \widehat{\theta}(i) < \pi, \text{ for all } i \geq 1, \text{ and } \lim_{i \rightarrow \infty} (\widehat{\theta}(i+1) - \widehat{\theta}(i)) = 0,$$

$$(A.4)$$

$$\lim_{r \rightarrow \infty} \sum_{i=1}^r (\widehat{\theta}(i+1) - \widehat{\theta}(i)) = \lim_{r \rightarrow \infty} \sum_{i=1}^r (\widehat{\rho}(i) - 1) = \lim_{r \rightarrow \infty} \sum_{i=1}^r \left(\frac{1}{\widehat{\rho}(1) - 1} + i - 1 \right)^{-1} = \infty,$$

where the equality $\widehat{\rho}(i) - 1 = \left(\frac{1}{\widehat{\rho}(1) - 1} + i - 1 \right)^{-1}$ can be proved by induction over i . Properties (A.2), (A.3), and (A.4) ensure that there exists a sequence $\{n_h \mid h \in \mathbb{Z}_{\geq 0}\}$ such that $(\rho(n_h), \theta(n_h)) = T_2(\rho(n_h - 1), \theta(n_h - 1))$ for all $h \in \mathbb{Z}_{\geq 0}$, and $(\rho(t), \theta(t)) = T_1(\rho(t - 1), \theta(t - 1))$ if $t \notin \{n_h \mid h \in \mathbb{Z}_{\geq 0}\}$. Moreover, we have that $\lim_{h \rightarrow \infty} (n_{h+1} - n_h) = \infty$. In other words, both the maps T_1 and T_2 are applied *infinitely often* along the evolution described by $\{(\rho(n), \theta(n))\}$, but there does not exist $D \in \mathbb{N}$ such that T_2 is applied at least once within each interval $[n, n + D]$, $n \in \mathbb{Z}_{\geq 0}$. Observe that, in this case, $(\rho(n), \theta(n))$ converges to the set $\{(1, \theta) \mid \theta \in [0, 2\pi]\} \subset X$. This set is different from the ω -limit set in equation (A.1).

Appendix B. Discontinuity of the multicenter function in the Hausdorff metric. To see that $\mathcal{H}_{\text{centroid}} : \mathcal{V}_N^* \rightarrow \mathbb{R}_{\geq 0}$ is not Hausdorff-continuous, consider the sequence of 2-partitions $\{v(t)\}_{t \in \mathbb{Z}_{\geq 0}}$ of the interval $[-1, 1] \subseteq \mathbb{R}$ defined by

$$v_1(t) = \left[-1, -1 + \frac{1}{2^{t+1}} \right] \cup \bigcup_{h=-(2^{t-1}-1)}^{2^{t-1}-1} \left[\frac{h}{2^{t-1}} - \frac{1}{2^{t+1}}, \frac{h}{2^{t-1}} + \frac{1}{2^{t+1}} \right] \cup \left[1 - \frac{1}{2^{t+1}}, 1 \right],$$

and by $v_2(t) = \overline{[-1, 1] \setminus v_1(t)}$. Note that both sequences $\{v_1(t)\}_{t \in \mathbb{Z}_{\geq 0}}$ and $\{v_2(t)\}_{t \in \mathbb{Z}_{\geq 0}}$ converge to $[-1, 1]$, and that $\text{Cd}(v_1(t)) = \text{Cd}(v_2(t)) = \text{Cd}([-1, 1]) = 0$, for all $t \geq 0$. Hence, for $\phi(s) = 1$ and $f(s) = s$, we compute

$$\mathcal{H}_1(0, v_1(t)) = \int_{v_1(t)} |x| dx = 2 \int_{v_1(t) \cap [0, 1]} |x| dx = 1,$$

and consequently $\mathcal{H}_{\text{centroid}}(v(t)) = 2$, while $\mathcal{H}_{\text{centroid}}(\lim_{t \rightarrow \infty} v(t)) = 2\mathcal{H}_1(0, [-1, 1]) = 4$. This shows that $\lim_{t \rightarrow \infty} \mathcal{H}_{\text{centroid}}(v(t)) \neq \mathcal{H}_{\text{centroid}}(\lim_{t \rightarrow \infty} v(t))$.

REFERENCES

- [1] E. S. ADAMS, *Approaches to the study of territory size and shape*, Annual Review of Ecology and Systematics, 32 (2001), pp. 277–303.
- [2] F. R. ADLER AND D. M. GORDON, *Optimization, conflict, and nonoverlapping foraging ranges in ants*, American Naturalist, 162 (2003), pp. 529–543.
- [3] P. K. AGARWAL AND M. SHARIR, *Efficient algorithms for geometric optimization*, ACM Computing Surveys, 30 (1998), pp. 412–458.
- [4] A. ARSIE, K. SAVLA, AND E. FRAZZOLI, *Efficient routing algorithms for multiple vehicles with no explicit communications*, IEEE Transactions on Automatic Control, 54 (2009), pp. 2302–2317.
- [5] A. BACCIOTTI AND L. MAZZI, *An invariance principle for nonlinear switched systems*, Systems & Control Letters, 54 (2005), pp. 1109–1119.
- [6] G. W. BARLOW, *Hexagonal territories*, Animal Behavior, 22 (1974), pp. 876–878.
- [7] S. BOYD, A. GHOSH, B. PRABHAKAR, AND D. SHAH, *Randomized gossip algorithms*, IEEE Transactions on Information Theory, 52 (2006), pp. 2508–2530.

- [8] R. W. BROCKETT, *System theory on group manifolds and coset spaces*, SIAM Journal on Control, 10 (1972), pp. 265–284.
- [9] F. BULLO, R. CARLI, AND P. FRASCA, *Gossip coverage control for robotic networks: Dynamical systems on the space of partitions*, Dec. 2009. Available at <http://arxiv.org/abs/0903.3642>.
- [10] F. BULLO, J. CORTÉS, AND S. MARTÍNEZ, *Distributed Control of Robotic Networks*, Applied Mathematics Series, Princeton University Press, 2009. Available at <http://www.coordinationbook.info>.
- [11] J. CORTÉS, S. MARTÍNEZ, T. KARATAS, AND F. BULLO, *Coverage control for mobile sensing networks*, IEEE Transactions on Robotics and Automation, 20 (2004), pp. 243–255.
- [12] M. DE BERG, M. VAN KREVELD, M. OVERMARS, AND O. SCHWARZKOPF, *Computational Geometry: Algorithms and Applications*, Springer, 2 ed., 2000.
- [13] Z. DREZNER AND H. W. HAMACHER, eds., *Facility Location: Applications and Theory*, Springer, 2001.
- [14] Q. DU, M. EMELIANENKO, AND L. JU, *Convergence of the Lloyd algorithm for computing centroidal Voronoi tessellations*, SIAM Journal on Numerical Analysis, 44 (2006), pp. 102–119.
- [15] Q. DU, V. FABER, AND M. GUNZBURGER, *Centroidal Voronoi tessellations: Applications and algorithms*, SIAM Review, 41 (1999), pp. 637–676.
- [16] Q. DU AND D. WANG, *Anisotropic centroidal Voronoi tessellations and their applications*, SIAM Journal on Computing, 26 (2005), pp. 737–761.
- [17] J. W. DURHAM, R. CARLI, P. FRASCA, AND F. BULLO, *Discrete partitioning and coverage control with gossip communication*, in ASME Dynamic Systems and Control Conference, Hollywood, CA, USA, Oct. 2009, pp. 225–232.
- [18] R. GRAHAM AND J. CORTÉS, *Asymptotic optimality of multicenter Voronoi configurations for random field estimation*, IEEE Transactions on Automatic Control, 54 (2009), pp. 153–158.
- [19] R. M. GRAY AND D. L. NEUHOFF, *Quantization*, IEEE Transactions on Information Theory, 44 (1998), pp. 2325–2383. Commemorative Issue 1948-1998.
- [20] H. GRÖMER, *On the symmetric difference metric for convex bodies*, Beiträge zur Algebra und Geometrie (Contributions to Algebra and Geometry), 41 (2000), pp. 107–114.
- [21] A. GUSRIALDI, S. HIRCHE, D. ASIKIN, T. HATANAKA, AND M. FUJITA, *Voronoi-based coverage control with anisotropic sensors and experimental case study*, Intelligent Service Robotics, 2 (2009), pp. 195–204.
- [22] J. HESPANHA, D. LIBERZON, D. ANGELI, AND E. D. SONTAG, *Nonlinear norm-observability notions and stability of switched systems*, IEEE Transactions on Automatic Control, 50 (2005), pp. 154–168.
- [23] A. K. JAIN, R. P. W. DUIN, AND J. MAO, *Statistical pattern recognition: A review*, IEEE Transactions on Pattern Analysis & Machine Intelligence, 22 (2000), pp. 4–37.
- [24] D. KEMPE, A. DOBRA, AND J. GEHRKE, *Gossip-based computation of aggregate information*, in IEEE Symposium on Foundations of Computer Science, Washington, DC, Oct. 2003, pp. 482–491.
- [25] S. G. KRANTZ AND H. R. PARKS, *Geometric Integration Theory*, Birkhäuser, 2008.
- [26] S. P. LLOYD, *Least squares quantization in PCM*, IEEE Transactions on Information Theory, 28 (1982), pp. 129–137. Presented at the 1957 Institute for Mathematical Statistics Meeting.
- [27] D. G. LUENBERGER, *Linear and Nonlinear Programming*, Addison-Wesley, 2 ed., 1984.
- [28] J. MACQUEEN, *Some methods for the classification and analysis of multivariate observations*, in Proceedings of the Fifth Berkeley Symposium on Mathematics, Statistics and Probability, University of California Press, 1965-1966, pp. 281–297.
- [29] S. MARTÍNEZ, F. BULLO, J. CORTÉS, AND E. FRAZZOLI, *On synchronous robotic networks – Part II: Time complexity of rendezvous and deployment algorithms*, IEEE Transactions on Automatic Control, 52 (2007), pp. 2214–2226.
- [30] A. MOSSER AND C. PACKER, *Group territoriality and the benefits of sociality in the African lion*, Panthera leo, Animal Behaviour, 78 (2009), pp. 359–370.
- [31] S. B. NADLER, *Hyperspaces of sets*, vol. 49 of Monographs and Textbooks in Pure and Applied Mathematics, Marcel Dekker, 1978.
- [32] M. PAVONE, A. ARSIE, E. FRAZZOLI, AND F. BULLO, *Distributed algorithms for environment partitioning in mobile robotic networks*, IEEE Transactions on Automatic Control, 56 (2011), pp. 1834–1848.
- [33] R. T. ROCKAFELLAR, *Convex Analysis*, Princeton University Press, 1970.
- [34] G. SALINETTI AND R. WETS, *On the convergence of sequences of convex sets in finite dimensions*, SIAM Review, 21 (1979), pp. 18–33.
- [35] R. G. SANFELICE, R. GOEBEL, AND A. R. TEEL, *Invariance principles for hybrid systems with*

- connections to detectability and asymptotic stability*, IEEE Transactions on Automatic Control, 52 (2007), pp. 2282–2297.
- [36] A. SARLETTE AND R. SEPULCHRE, *Consensus optimization on manifolds*, SIAM Journal on Control and Optimization, 48 (2009), pp. 56–76.
- [37] M. SCHWAGER, D. RUS, AND J. J. SLOTINE, *Decentralized, adaptive coverage control for networked robots*, International Journal of Robotics Research, 28 (2009), pp. 357–375.
- [38] S. SUSCA, S. MARTÍNEZ, AND F. BULLO, *Monitoring environmental boundaries with a robotic sensor network*, IEEE Transactions on Control Systems Technology, 16 (2008), pp. 288–296.
- [39] M. TANEMURA AND M. HASEGAWA, *Geometrical models of territory. I. Models for synchronous and asynchronous settlement of territories*, Journal on Theoretical Biology, 82 (1980), pp. 477–496.
- [40] J. N. TSITSIKLIS, D. P. BERTSEKAS, AND M. ATHANS, *Distributed asynchronous deterministic and stochastic gradient optimization algorithms*, IEEE Transactions on Automatic Control, 31 (1986), pp. 803–812.
- [41] R. VOTEL, D. A. W. BARTON, T. GOTOU, T. HATANAKA, M. FUJITA, AND J. MOEHLIS, *Equilibrium configurations for a territorial model*, SIAM Journal on Applied Dynamical Systems, 8 (2009), pp. 1234–1260.